



Preliminary Environmental Information Report

Volume III - Appendices

Appendix 14B: Fisheries and Fish Ecology Baseline

The Infrastructure Planning (Environmental Impact Assessment) Regulations 2017 (as amended)



Prepared by: **AECOM**

Table of Contents

14B. Fisheries and Fish Ecology Baseline	14-4
14.1 Introduction.....	14-4
14.2 Methodology.....	14-5
14.3 Marine Designations.....	14-8
14.4 Fish and Shellfish Communities	14-11
14.5 Commercial Fisheries.....	14-24
14.6 Species-Specific Information	14-32
14.7 Baseline Evolution	14-53
14.8 Summary of Findings.....	14-55
14.9 References	14-57
Annex A – Temporal variations in the type of gear used during the Environmental TraC surveys	14-68
Annex B – Full species list and proportions as presented in TraC survey fish species proportion figures.	14-72
Annex C – Freshwater Fish Count NFPD Environment Agency Survey.....	14-75
Figures	14-76

Diagrams

Diagram 14B-1: Diadromous fish counts from Environment Agency (2019a) TraC surveys in the River Tees. Data has been reported for a range of survey methods (beam trawl, otter trawl, and seine netting) and effort.	14-17
Diagram 14B-2: Total fish counts for each gear type used, in Environment Agency (2019a) TraC surveys in the lower reaches of the River Tees.	14-19
Diagram 14B-3: Proportion of fish species from Environment Agency (2019a) TraC surveys in the lower reaches of the River Tees (1985 – 2018). Fish species in the legend have been presented in order of percentage contribution.	14-20
Diagram 14B-4: Total fish counts for each gear type used, in Environment Agency (2019a) TraC surveys in the middle reaches of the River Tees.	14-21
Diagram 14B-5: Proportion of fish species from Environment Agency (2019a) TraC surveys in the middle reaches of the River Tees (1981 – 2015). Fish species in the legend have been presented in order of percentage contribution.	14-21
Diagram 14B-6: Total fish counts for each gear type used, in Environment Agency (2019a) TraC surveys in the upper reaches of the River Tees.	14-22
Diagram 14B-7: Proportion of fish species from Environment Agency (2019) TraC surveys in the upper reaches of the River Tees (1982 – 1996). Fish species in the legend have been presented in order of percentage contribution.	14-23
Diagram 14B-8: Average landed weight (tonnes) (2013 – 2017) recorded in the ICES rectangle 38E8 for each gear category used. Fishing activity data taken from iFISH data system, as reported by the MMO (2018).	14-27

Diagram 14B-9: Annual landed weight (tonnes) (2013 – 2017) recorded in the ICES rectangle 38E8. Fishing activity data taken from IFISH data system, as reported by the MMO (2018).	14-28
Diagram 14B-10: Average landed weight (tonnes) (2013 – 2017) recorded in the ICES rectangle 38E8 for each vessel length category. Fishing activity data taken from IFISH data system, as reported by the MMO (2018).	14-31
Diagram 14B-11: Main salmon rivers in England and Wales, denoting those with Salmon Action Plans (*) and those designated as Special Areas of Conservation (\$) in which salmon must be maintained or restored to favourable conservation status (CEFAS <i>et al.</i> , 2019).	14-34
Diagram 14B-12: Monthly combined upstream counts for salmon and sea trout from 2011 to 2019 at the Tees Barrage on the lower Tees, reported by the Environment Agency (Environment Agency, 2019b). Counter only monitors upstream migration through the fish pass. No data was reported for January to June 2011 and August to December 2019.	14-35
Diagram 14B-13: Monthly mean upstream counts for salmon from 1995 to 2011 at the Tees Barrage on the lower Tees, reported by the Environment Agency (Environment Agency, 2013). Fish trap counts only refer to upstream migration through the fish pass. Error bars refer to standard deviation. No data was reported for January to April 1995 and July to December 2011.	14-36
Diagram 14B-14: Monthly mean upstream counts for sea trout from 1995 to 2011 at the Tees Barrage on the lower Tees, reported by the Environment Agency (Environment Agency, 2013). Fish trap counts only refer to upstream migration through the fish pass. Error bars refer to standard deviation. No data was reported for January to April 1995 and July to December 2011.	14-36
Diagram 14B-15: Summary of salmon and sea trout rod catches from the river Tees that were declared from 2007 to 2017, as reported by the Environment Agency (2017).	14-37
Diagram 14B-16: Total fish counts for European eel (<i>Anguilla anguilla</i>), in Environment Agency (2019a) TraC surveys in the three reaches of the River Tees.	14-39
Diagram 14B-17: UK Distribution of River Lamprey (left) and Sea Lamprey (right) (JNCC, 2013a and 2013b).	14-40
Diagram 14B-18: Size classes (mm) (based on total length (mm)) measured for herring (<i>Clupea harengus</i>) in Environment Agency (2019a) TraC surveys in the River Tees. Data has been presented for the first 95% of fish counts.	14-42
Diagram 14B-19: Size classes (mm) (based on total length (mm)) measured for sprat (<i>Sprattus sprattus</i>) in Environment Agency (2019a) TraC surveys in the River Tees. Data has been presented for the first 95% of fish counts.	14-43
Diagram 14B-20: Size classes (mm) (based on total length (mm)) measured for cod (<i>Gadus morhua</i>) in Environment Agency (2019a) TraC surveys in the River Tees. Data has been presented for the first 95% of fish counts.	14-46
Diagram 14B-21: Size classes (mm) (based on total length (mm)) measured for whiting (<i>Merlangus merlangus</i>) in Environment Agency (2019a) TraC surveys in the River Tees. Data has been presented for the first 95% of fish counts.	14-47
Diagram 14B-22: Size classes (mm) (based on total length (mm)) measured for plaice (<i>Pleuronectes platessa</i>) in Environment Agency (2019a) TraC surveys in the River Tees. Data has been presented for the first 95% of fish counts.	14-49

Diagram 14B-23: Size classes (mm) (based on total length (mm)) measured for *Ammodytidae* (sandeel species) in Environment Agency (2019a) TraC surveys in the River Tees. Data has been presented for the first 95% of fish counts. 14-51

Tables

Table 14B-1: Summary of key legislation relevant to the assessment of fisheries and fish ecology baseline	14-5
Table 14B-2: Summary of relevant fish and shellfish species protected by national and international legislation or policy.....	14-10
Table 14B-3: Marine fish counts from Environment Agency (2019a) TraC surveys in the River Tees for 2009 to 2018. Data has been reported for a range of survey methods (beam trawl, otter trawl, and seine netting) and effort. Fish species have been presented in order of prevalence (number of years present), followed by total abundance.	14-15
Table 14B-4: Nationalities of vessels fishing in the ICES rectangle 38E8 and the weight (tonnes) of their annual landings (tonnes) (2013 – 2017).	14-24
Table 14B-5: Length categories of vessels fishing in the ICES rectangle 38E8 and their average landed weight (tonnes) per year (2013 – 2017) for each gear type used.	14-29

14B. Fisheries and Fish Ecology Baseline

14.1 Introduction

Aims and Objectives

- 14.1.1 This study aims to provide a detailed baseline description of marine and estuarine fish and shellfish populations (including diadromous species) and fisheries within the Tees Estuary and the wider coastal marine environment in the absence of the Proposed Development. The objective is to enable identification of potentially important fisheries and fish ecology features within the Zone of Influence (Zol) of the Proposed Development.
- 14.1.2 This report presents the results of a desk study carried out to date, the objectives of which were to:
- Collate and review available information on fisheries and fish ecology;
 - Summarise the findings of the available information; and
 - Identify key fish and shellfish receptors as well as relevant marine designations in the area.
- 14.1.3 No project-specific fish surveys have been undertaken or are proposed.

Study Area

- 14.1.4 Based on the location of the Proposed Development, the study area for the fisheries and fish ecology baseline has been defined as the area comprising the River Tees, the Tees estuary, and the wider coastal area up to and including the Greater North Sea out to a distance of 10 km offshore from the indicative DCO boundary (see Figure 14B-1, PEI Report, Volume III). This spatial extent was chosen on the basis that it provides geographical context and encompasses the relevant functional habitats and range of movement for the species found within the vicinity of the Proposed Development.
- 14.1.5 The study area falls within the MMO North East Inshore Marine Plan area and the International Council for the Exploration of the Sea (ICES) rectangle 38E8, representing a standard geographical unit, of approximately 30 by 30 nautical miles, for the reporting of fisheries data. In addition, the study area is also encompassed within the district of the North Eastern Inshore Fisheries and Conservation Authority (NEIFCA) which is responsible for managing and conserving marine resources between the River Tyne and North East Lincolnshire.

Structure of Report

- 14.1.6 This report is structured as follows:

- **Section 14.2 (Methodology)** – summarises the methodology for undertaking the fisheries and fish ecology baseline study and includes information related to relevant conservation legislation and key data sources;
- **Section 14.3 (Marine Designations)** – provides an overview of marine designations including local, national and internationally protected sites and species which are relevant to the assessment of the fisheries and fish ecology baseline;
- **Section 14.4: (Fish and Shellfish Communities)** – provides a general characterisation of fish and shellfish communities within the study area and presents the results of the National Fish Populations Database (NFPD) for the Tees as reported by the Environment Agency (2019a);
- **Section 14.5: (Commercial Fisheries)** – describes the commercial fish and shellfish fisheries operating within the study area;
- **Section 14.6: (Species Information)** – provides further information on the ecology of specific fish and shellfish species known to be of commercial and/or conservational importance in the study area;
- **Section 14.7 (Baseline Evolution)** – summarises how the fisheries and fish ecology baseline may change during the consenting process and over the lifetime of the Proposed Development;
- **Section 14.8 (Summary of Findings)** – provides a summary of the findings of the desk study and identifies the key fish and shellfish receptors which require consideration within the relevant environmental assessments.

14.2 Methodology

Legislative Context

- 14.2.1 Key biodiversity and conservation legislation relevant to the development of the fisheries and fish ecology baseline is shown in Table 14B-1. This information has been used to inform the scope of baseline information required and to determine the key ecological features pertinent to the assessment of effects outlined within the Environmental Statement (ES).

Table 14B-1: Summary of key legislation relevant to the assessment of fisheries and fish ecology baseline

Name	Description	How it relates to the Proposed Development
The Marine Strategy Regulations 2010	Transposes the Marine Strategy Framework Directive (MSFD) (2008/56/EC) into UK legislation. The MSFD sets out a framework for achieving Good Environmental Status by 2020. As such, member states are required to carry out an assessment of the current state of UK marine waters and establish	The latest UK assessment states that GES is not currently being achieved for fish (including some commercial fish and shellfish species). The four ecology indicators used to make this assessment include: population abundance, size structure and species composition. For commercial species, indicators

Name	Description	How it relates to the Proposed Development
	what constitutes good environmental status, including the development of relevant targets and indicators.	include commercial fishing pressure and reproductive capacity. These indicators have been considered as part of this baseline characterisation.
The Water Environment (Water Framework Directive (WFD)) Regulations 2017	Transposes the European Water Framework Directive (2000/60/EC) in UK legislation. It establishes a legislative framework for the protection of surface waters (including rivers, lakes, transitional waters and coastal waters) and groundwater.	Fish represents a biological quality element considered in the determination of ecological status for the Tees and the Tees Coastal water bodies which fall within the study area.
The Conservation of Habitats and Species Regulations 2017 ("The Regulations")	The Regulations transpose the Habitats Directive (92/43/EEC) into UK legislation out to the 12 nautical mile (nm) limit and provide for the designation and protected of European Sites, the protection of 'European protected species', and the adaption of planning and other controls for the protection of European Sites.	There are several European Sites and European protected fish species listed under Annex II and IV known to be present within the study area and require consideration.
Section 41 of the Natural Environment and Rural Communities (NERC) Act 2006 (England)	Provides a statutory list of species and habitats of principal importance for the purpose of conservation biodiversity.	Several species known to be present within the study area are listed under Section 41 as being of Principal Importance in England.
The Convention on the Conservation of Migratory Species of Wild Animals (the 'Bonn Convention') 1985	Provides protection for endangered migratory species (listed in Appendix I and II) and their habitats. The legal requirement for the strict protection of Appendix I & II species is provided by the Wildlife & Countryside Act (1981 as amended).	Species listed in Appendix II are known to present within the study area.
The Convention on the Conservation of European Wildlife and Natural Habitats (the Bern Convention) 1982	Aims to conserve and protect wild plant and animal species and their natural habitats (listed in Appendices II and III of the Convention). The legal requirement for the strict protection of Appendix II & III species is provided by the Wildlife & Countryside Act (1981 as amended).	Species listed in Appendix III are known to be present within the study area.
The Oslo and Paris Convention for the Protection of the Marine Environment of the North-East Atlantic (the 'OSPAR Convention') 1998	Initiated the development of a list of threatened and/or declining species and habitats that required protection in accordance OSPAR Biological Diversity and Ecosystem Strategy.	Species listed by OSPAR are known to be present within the study area.
Convention on Biological Diversity 1992	Provides an international legal framework for the conservation of biodiversity and sustainable use of	Highlights the importance of conserving, protecting and enhancing general biological

Name	Description	How it relates to the Proposed Development
	its components. UK Biodiversity Action Plan (UK BAP)* priority species and habitats lists were generated in response to this piece of legislation.	diversity (as well as specific species and habitats). Several UK BAP priority fish species are known to be present within the study area.

* The 'UK Post-2010 Biodiversity Framework', was published in 2012 and succeeds the UK BAP (Joint Nature Conservation Committee (JNCC) and Defra, 2018). Despite this, the UK BAP list of priority species and habitats remains an important reference material.

14.2.2 In England, fisheries management between 0 – 6 nautical miles (nm) is under the jurisdiction of the Inshore Fisheries Conservation Authorities (IFCAs), which were established under the Marine and Coastal Access Act (MCAA) 2010. The IFCAs main legal duties are described in Section 153 of the MCAA and they have a duty to sustainably manage sea fisheries resources within their district and to protect marine ecosystems from the impact of fishing. As outlined in paragraph 14.1.5, the study area falls within the North Eastern IFCA (NEIFCA) District.

14.2.3 The Environment Agency has a general duty to maintain, improve and develop freshwater fisheries of migratory species, including salmon, trout and freshwater eels. The Salmon and Freshwater Fisheries Act 1975 (as amended) provides the Agency with the powers to make Byelaws and restrict the number of licences issued for individual net fisheries by making Net Limitation Orders (NLO's).

Data Sources

14.2.4 The fisheries and fish ecology baseline have been described using several data sources. These data sources were used to determine the relative importance and functionality of the study area in the regional context of fish populations in the Teesside estuarine/coastal area and the associated North Sea, as well as the populations of diadromous fish using the River Tees. The data sources reviewed include the following:

- General:
 - FishBase (www.fishbase.org) for general fish ecology, distribution and biological information;
 - The Humber and East Coast Regional Environmental Characterisation (Tappin *et al.*, 2011 and Limpenny *et al.*, 2011, respectively) for a summary of the distribution and ecology of fish in the North Sea;
 - Environment Agency (2009) River Tees Salmon Action Plan which sets out stock assessments and management actions for this species;
 - Centre for Environment, Fisheries and Aquaculture (CEFAS) Sensitivity Maps (Coull *et al.*, 1998; Ellis *et al.*, 2012) which provide spatial data highlighting spawning and nursery grounds in UK waters;

- The International Convention for the Conservation of Nature (IUCN) Red List of Threatened Species (<https://www.iucnredlist.org/>); and
- Published Environmental Statements and survey reports produced for large infrastructure projects in the vicinity of the Proposed Development, including Dogger Bank Teesside A & B Offshore Wind (Brown and May Marine, 2014; Precision Marine Survey Ltd, 2014) and Teesside Offshore Wind Farm (Lancaster *et al.*, 2011).
- Data:
 - Environment Agency (2019a) National Fish Populations Database Transitional and Coastal (TraC) fish counts for all species within the Tees Estuary;
 - Environment Agency (2013, 2019b) fish count data recorded from the Tees Barrage;
 - Environment Agency (2019c) National Fish Populations Database freshwater fish counts for species within the River Tees
 - Marine Mammal Organisation (MMO) 2013 – 2014 annual landings statistics (<https://www.gov.uk/government/collections/uk-sea-fisheries-annual-statistics>);
 - Environment Agency (2017). Salmonid and Freshwater Fisheries Statistics for England and Wales 2017;
 - CEFAS (2019) assessment of salmon stock and fisheries in England and Wales; and
 - International Council for the Exploration of the Seas (ICES) data (<https://www.ices.dk/Pages/default.aspx>).

14.3 Marine Designations

Relevant Marine Designated Sites

- 14.3.1 To the north east of the Main Site, as part of the Proposed Development, lie the coastal areas of South Gare and Cotham Sands which form part of the Teesmouth and Cleveland Coast Special Protected Area (SPA) / Ramsar site. Underpinning the SPA designation is the Teesmouth and Cleveland Coast Site of Specific Interest (SSSI), which encompasses five SSSI sites within 5 km of the indicative DCO site boundary (see Figure 14B-2, PEI Report, Volume III). Both the Teesmouth and Cleveland Coast SPA and SSSI are designated for the protected breeding / non-breeding bird species and other important waterfowl species associated with the site. As of January 2020, the proposed extension of the existing Teesmouth and Cleveland SPA and Ramsar site has been formally adopted and is intended to protect important marine foraging areas for breeding terns well as intertidal areas and estuarine waters used by wintering birds.
- 14.3.2 The Teesmouth and Cleveland Coast SPA / Ramsar site includes a range of coastal habitats (sand-flats and mud-flats, rocky shore, saltmarsh, freshwater marsh and sand dunes) within and around the Tees Estuary. These habitats are important in providing feeding and roosting opportunities

for important numbers of water birds in winter and during passage periods. This includes in summer when Little Tern (*Sterna albifrons*) breed on the beaches within the site and when Sandwich Tern (*Sterna sandvicensis*) are abundant on passage. These birds, as well as others also protected under the site designation are known to prey upon small fish species including juvenile sandeel (*Ammodytes* spp.), clupeids such as herring (*Clupea harengus*) and sprat (*Sprattus sprattus*), and sand smelt (*Atherina presbyter*), as well as molluscs, crustaceans and worms (Green, 2017).

- 14.3.3 The Teesmouth and Cleveland Coast SSSI is of special interest for a range of nationally important features and supported by a wider mosaic of coastal and freshwater habitats. The SSSI includes the whole of the Tees Estuary, from between North Gare and South Gare, where the mouth of the estuary lies, up to the tidal limits of the Tees and Greatham Creek. As with the SPA, the key features of the SSSI include both non-breeding and breeding birds which utilise and assemble on the sand dune and saltmarsh special features and lowland open waters and are known to prey on small fish species.
- 14.3.4 Seal Sands SSSI, located approximately 2.9 km to the west of the DCO boundary, supports a population of harbour seal (*Phoca vitulina*) which re-colonised the area in the 1980s and has since established a regular breeding colony. Harbour seals in the British Isles are known to prey on a variety of fish species including: sandeel, gadoids, flatfish, herring and sprat; the large majority consumed being <30 cm in estimated length (Wilson and Hammond, 2016). Grey seals are also known to be present in the study area and prey upon similar fish species to harbour porpoise (SCOS, 2018).
- 14.3.5 There are no other European Sites or Marine Conservation Zones within 10 km of the indicative DCO boundary.

Designated Species

- 14.3.6 The study area of the Proposed Development is found within the Tees Valley BAP which covers the local authority areas of Hartlepool, Stockton-on-Tees, Middlesbrough, and Redcar and Cleveland. A variety of migratory fish species, which are species of Principal Importance in England, have been identified as utilising the Tees Estuary and forming the basis of the local BAP. These species include salmon (*Salmo salar*), sea trout (*Salmo trutta*), European eel (*Anguilla anguilla*), sea lamprey (*Petromyzon marinus*), and river lamprey (*Lampetra fluviatilis*). Furthermore, the River Tees is also recognised as a main salmon river in England and Wales, with a Salmon Action Plan (Environment Agency, 2009) enforced by the Environment Agency, used to provide a strategy for the management of the fishery.
- 14.3.7 Table 14B-2 lists all the fish known to be present in the study area which are protected under national and international conservation legislation. All species listed are also considered to be of commercial importance within the study area with the exception of sandeel and the diadromous fish species.
- 14.3.8 There are no shellfish species which are afforded conservation protection known to be present in the study area.

Table 14B-2: Summary of relevant fish and shellfish species protected by national and international legislation or policy.

Common names	Latin names	Habitats Directive Annex II and IV species	OSPAR list of threatened and/or declining species	Bonn Convention Appendix I and II species	Bern Convention Appendix II and III species	UK BAP Priority Species	NERC 2006 Species of Principal Importance	Features of Conservation Interest (FOCI)	IUCN Red List*
Pelagic fish species									
Herring	<i>Clupea harengus</i>					✓	✓	✓	LC (↑)
Mackerel	<i>Scomber scombrus</i>					✓	✓	✓	LC (↓)
Demersal fish species									
Cod	<i>Gadus morhua</i>		✓			✓	✓	✓	VU (-)
Whiting	<i>Merlangius merlangus</i>					✓	✓	✓	LC (?)
Dover sole	<i>Solea solea</i>					✓	✓	✓	DD (↔)
Plaice	<i>Pleuronectes platessa</i>					✓	✓	✓	LC (↑)
Sandeel	<i>Ammodytidae</i>					✓ ¹	✓ ¹	✓ ¹	LC (?) ¹
Diadromous fish species									
Atlantic salmon	<i>Salmo salar</i>	✓	✓			✓	✓	✓	LC (-)
Sea trout	<i>Salmo trutta</i>					✓	✓		LC (?)
European eel	<i>Anguilla anguilla</i>		✓	✓		✓	✓	✓	CR (↓)
Sea lamprey	<i>Petromyzon marinus</i>	✓	✓		✓	✓	✓	✓	LC (↔)
River lamprey	<i>Lampetra fluviatilis</i>	✓				✓	✓	✓	LC (?)

* IUCN Red List Status defined as 'CR' = Critically Endangered, 'VU' = Vulnerable, 'NT' = Near Threatened, 'LC' = Least Concern and 'DD' = Data Deficient. Population trends are also shown in brackets ('↑' = increasing, '↓' = decreasing, '↔' = stable, '?' = unknown and '-' = unspecified).

¹ Raitts sandeel (*Ammodytes marinus*) species.

² Norway lobster (*Nephrops norvegicus*) species.

14.4 Fish and Shellfish Communities

- 14.4.1 The central North Sea area is described as having a mixed demersal and pelagic fish assemblage with a high diversity of species, likely due to an overlap of the northern and southern North Sea fish communities (Reiss *et al.*, 2009). The fish assemblage in the deeper northern region of the North Sea (central and northern areas) is characterised by the following fish species: herring, mackerel (*Scomber scombrus*), horse mackerel (*Trachurus trachurus*), cod (*Gadus morhua*), whiting (*Merlangius merlangus*), haddock (*Melanogrammus aeglefinus*), plaice (*Pleuronectes platessa*), and dab (*Limanda limanda*) (Teal, 2011; Callaway *et al.*, 2002). Important prey fish species for the North Sea also include sandeels (Ammodytidae), Norway pout (*Trisopterus esmarki*), and sprat (ICES, 2008).
- 14.4.2 The occurrence, distribution and abundance of many fish and shellfish within the study area is determined by their propensity to aggregate within coastal areas to spawn (lay or release their eggs). ‘Spawning grounds’ are defined either by the species behaviour and therefore may cover a wide area, or by specific habitat preferences (e.g. gravel), which may restrict spatial extent. Fish exhibit several modes of reproduction, the most common being broadcast spawning, where eggs and sperm are released into the water column (Ellis *et al.*, 2012). Other species deposit egg-cases (e.g. dogfish and whelk) or egg mats onto the seafloor (e.g. herring), making them particularly vulnerable to seabed disturbance. Eggs which are fertilised in the water column (i.e. pelagic) are transported along with pelagic early life stages (e.g. larvae and juvenile fish) within the plankton to nursery areas which provide plentiful food and shelter for young fish species. Once grown, most fish leave their nursery grounds to join adult populations further from shore.
- 14.4.3 Fisheries sensitivity maps (Coull *et al.*, 1998; Ellis *et al.*, 2012) provide information on spawning grounds (the location where eggs are laid) and nursery areas (the location where juveniles are common) for selected fish and shellfish species prevalent in the study area. This data indicates that the Proposed Development is located within the nursery grounds of the following species: herring, anglerfish, plaice, cod, whiting and spurdog (*Squalus acanthias*) (Figure 14B-3, PEI Report, Volume III) as well as sprat, *nephrops* (Norway lobster) and lemon sole. The Proposed Development is also found within the spawning area of plaice, lemon sole and *nephrops*. Juvenile horse mackerel appear to be widespread exhibiting no spatially discrete nursery grounds within the study area.
- 14.4.4 Diadromous fish are known to transit through the River Tees and estuary during seasonal migrations between the sea and riverine environments. The species known to migrate through the study area were identified as part of the Tees Valley BAP and include salmon, sea trout, European eel, river lamprey and sea lamprey, all of which are UK priority BAP species (see Table 14B-2). The Tees river is particularly important for salmon and sea trout with these species regularly using the Tees catchment area (Moore and Potter, 2014). The River Tees includes a tidal barrage which was constructed across the Tees at Stockton in 1995 (Environment Agency, 2009). During a

three-year tracking study of fish passage at the River Tees Barrage, conducted by Moore and Potter (2014), a total of 237 fish, of which 84% were salmon, were recorded. Salmon and trout in the Tees form the basis of small net and rod recreational fisheries for the catchment which places pressure on the population in combination with other factors such as water quality and quantity, obstructions to migration, loss of spawning habitat, predation and climate change (Environment Agency, 2009).

- 14.4.5 Fish surveys were undertaken in 2010 as part of the Teesside Offshore Wind pre-construction baseline monitoring programme with several sampling locations overlapping with the study area for the Proposed Development (Entec UK Limited, 2011). Three fishing techniques were employed, including trawling, gill nets, and potting. A total of 31 fish species were caught, with catches dominated by whiting, whilst plaice, haddock, dab and Atlantic cod were also caught in high abundances. Species such as lemon sole, saithe (*Pollachius virens*) and herring were recorded but in lower abundances. The shellfish catches were dominated by edible crab (*Cancer pagurus*), squid (*Loligo vulgaris*), harbour crab (*Liocarcinus depurator*) and European lobster (*Homarus gammarus*). Of the fish and shellfish species observed, edible crab, European lobster and velvet swimming crab (*Necora puber*), as well as plaice and dab, were mostly captured inshore within the windfarm site. In contrast, species such as cod and *nephrops* were caught offshore, whilst Dover sole and lemon sole were caught inshore during the summer months and offshore in the winter months.
- 14.4.6 A similar community composition was also observed in nearshore fish and shellfish surveys undertaken in autumn 2012 and spring 2013 to inform the Dogger Bank Teesside A & B export cable corridor ES (Precision Marine Survey Ltd, 2014). The fish surveys utilised the methods of otter trawling, scientific beam trawls and static gear (nets and shellfish pots). The results of these surveys found that, similar to the Teesside Offshore Wind surveys, whiting dominated catches, representing 47% of the total across all sites and surveys. Pouting (*Trisopterus luscus*), haddock, dab, plaice, and grey gurnard (*Eutrigla gurnardus*) were also caught in higher abundances. From the shellfish surveys, edible crab represented the highest proportion of the catch, contributing 48% to the overall total. Hermit crab (*Pagurus bernhardus*), star fish (*Asterias rubens*), whelk (*Buccinum undatum*), velvet swimmer crabs and lobster were also caught in relatively high abundance, representing 19%, 10%, 9%, 7%, and 5% of the total, respectively.

NFPD Fish Counts - Environment Agency

- 14.4.7 Data from the National Fish Populations Database (NFPD) for the Tees area, as reported by the Environment Agency (2019a; and 2019c), has been analysed and presented below. The NFPD provides a collection of information from fisheries monitoring work on rivers, lakes and transitional and coastal waters (TraC), recorded by the Environment Agency and third parties. The data includes information on fish counts, lengths, age and weight for a variety of marine, diadromous and freshwater fish species in different reaches of the River Tees.

14.4.8 It should be noted that the surveys represent a range of gear types and survey effort, meaning that the comparison of abundances is only indicative. The gear types used as part of the TraC fish surveys include:

- beam trawl – 1.5 m wide;
- beam trawl – 2.4 m wide;
- otter trawl netting; and
- seine netting.

14.4.9 A table summarising temporal and seasonal variations in the type of gear used in each area of the River Tees as part of the TraC fish surveys, can be found in Annex A. Typically, beam trawling with an aperture of 2.4 m was used from the 1980s to 2004 in both the middle and lower reaches and was the only technique used in the lower reaches across all years. From 2004 onwards, seine netting was used more frequently in the middle and lower reaches, whilst otter trawling was utilised only occasionally. Beam trawls with an aperture of 1.5 m were only used in the lower reaches of the river in all years from 2011.

14.4.10 In addition to the TraC fish surveys, freshwater fish count NFPD surveys were undertaken by the Environment Agency from 1993 to 2019 in the River Tees and associated tributaries. These surveys comprised of the following methodologies:

- electric fishing;
- seine netting; and
- fixed trap fishing.

14.4.11 The information from these surveys has been used as supporting evidence for the abundances and spatial distribution of the diadromous fish species present in the freshwater sections of the River Tees and its adjoining tributaries. As with the TraC surveys, these surveys represent a range of survey methodologies and survey effort, meaning that the comparison of abundances is only indicative.

Estuarine and Marine Fish Species

14.4.12 Table 14B-3 shows the marine fish species recorded across all areas of the River Tees in the most recent 10-year sampling period of the TraC monitoring surveys (2009 – 2018). It should be noted that monitoring spanned 1981 to 2018 but monitoring has focussed on the lower reaches only in the last 10 years and so this data was considered the most geographically and temporally relevant to the assessment of estuarine and marine fish species.

14.4.13 The species which were most prevalent and found across all 10 monitoring years were herring, plaice and sprat. These species contributed 73% to the total number of estuarine or marine fish observed. Lesser sandeel (*Ammodytes tobianus*) and flounder were also found to be prevalent within the study area, occurring in nine out of the 10 monitoring years and contributing a further 15% to the total abundance. In addition to the

mentioned species, sand goby and whiting were also prevalent for most of the monitoring period, being recorded in nine of the 10 most recent monitoring years but occurred in comparatively lower abundances (Table 14B-3).

- 14.4.14 Of the least prevalent species, dragonet (*Callionymus lyra*), greater sandeel (*Hyperoplus lanceolatus*), gurnard species (*Triglidae*), lemon sole (*Microstomus kitt*), painted goby (*Pomatoschistus pictus*), sand smelt (*Atherina presbyter*), sea bass (*Dicentrarchus labrax*), and thornback ray (*Raja clavata*) were only recorded in one year, with the four latter species only having one individual recorded, each. It is possible that some of these species, such as sea bass, were under-sampled due to the survey techniques utilised, although other species, such as turbot, are less common generally and therefore lower frequency occurrence would be expected.

Diadromous Fish Species

- 14.4.15 Diagram 14B-1 shows the diadromous fish species recorded across all areas of the River Tees for all years (1981 – 2018) reported as part of the TraC monitoring surveys. Data has been presented for all years in order to encompass sampling undertaken in the upper reaches of the River Tees, for which fish abundances were only reported from 1982 to 1996.
- 14.4.16 European eels are the most prevalent species recorded, having been reported in a total of 16 years, the most recently being in 2008. This species has the highest recorded relative abundance of 178 individuals, including two elvers reported in 2008. Sea trout was the diadromous species most recently recorded, reported to be present in 2015, having first been recorded in 1988. Atlantic salmon has only been recorded in July 1998, with one individual being recorded as part of beam trawl (2.4 m) surveys in the middle reaches of the river. Similarly, river lamprey has been reported on only one occasion in 1992, with three individuals recorded in a beam trawl (2.4 m).

Table 14B-3: Marine fish counts from Environment Agency (2019a) TraC surveys in the River Tees for 2009 to 2018. Data has been reported for a range of survey methods (beam trawl, otter trawl, and seine netting) and effort. Fish species have been presented in order of prevalence (number of years present), followed by total abundance.

Common Name	Latin Name	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018
Sprat	<i>Sprattus sprattus</i>	456	4,741	655	1,639	310	54	7	276	7	51
Herring	<i>Clupea harengus</i>	1,647	495	42	378	639	1,619	459	29	10	566
Plaice	<i>Pleuronectes platessa</i>	250	12	16	27	330	236	396	246	103	31
Lesser sandeel	<i>Ammodytes tobianus</i>		58	933	120	158	114	206	1	373	19
Flounder	<i>Platichthys flesus</i>	40	153	471	185	127	10	143	5	21	
Sand goby	<i>Pomatoschistus minutus</i>	35	65	15	5	38	11	73	45	54	
Whiting	<i>Merlangius merlangus</i>	228	2		2	1	7	17	8	18	14
Dab	<i>Limanda limanda</i>	260				10	69	102	189	168	
Common goby	<i>Pomatoschistus microps</i>		1	25	4	42	21	57			
Cod	<i>Gadus morhua</i>	287			3		5		32	20	
Lesser weever	<i>Echiichthys vipera</i>	8		2	1			4	2		
Pogge	<i>Agonus cataphractus</i>	13						10	5	16	
Dover sole	<i>Solea solea</i>	2				1		1	1		
Corbin's sandeel	<i>Hyperoplus immaculatus</i>	69		1	1						
Nilsson's pipefish	<i>Syngnathus rostellatus</i>						6	1	2		
Pollack	<i>Pollachius virens</i>	441					1				
Long-spined sea scorpion	<i>Taurulus bubalis</i>	10							2		
5-bearded rockling	<i>Ciliata mustela</i>	3						1			

Common Name	Latin Name	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018
Greater pipefish	<i>Syngnathus acus</i>									1	2
Snake pipefish	<i>Entelurus aequoreus</i>	1								1	
Turbot	<i>Psetta maxima</i>						1			1	
Viviparous blenny	<i>Zoarces viviparus</i>			1		1					
Dragonet	<i>Callionymus lyra</i>	4									
Greater sandeel	<i>Hyperoplus lanceolatus</i>					3					
Gurnard sp.	<i>Triglidae</i>	3									
Lemon Sole	<i>Microstomus kitt</i>	2									
Painted goby	<i>Pomatoschistus pictus</i>						1				
Sand smelt	<i>Atherina presbyter</i>		1								
Sea bass	<i>Dicentrarchus labrax</i>	1									
Thornback ray	<i>Raja clavata</i>									1	

Source: Environment Agency, 2019a




Diagram 14B-1: Diadromous fish counts from Environment Agency (2019a) TraC surveys in the River Tees. Data has been reported for a range of survey methods (beam trawl, otter trawl, and seine netting) and effort.

14.4.17 Freshwater fish counts from NFPD Environment Agency surveys recorded the following diadromous species: Atlantic salmon, brown / sea trout, European eel, and lamprey (individuals of lamprey were not identified to species level). Of these species, brown / sea trout was recorded in the greatest abundances, with a total of 18,535 individuals counted from the River Tees and its connected tributaries between 1994 and 2019. Atlantic salmon was the second most abundant, with 11,528 individuals across all sites and sampling years. These species are also the most consistently recorded, being present in all years sampled². A total of 1,385 European eel and 215 lampreys have been recorded to date; both taxa were recorded for the first time in 2001 but have only been consistently recorded from 2011 onwards when sampling at, and in the vicinity of, the Tees Barrage began.

Spatial Variations

14.4.18 Figure 14B-4 (PEI Report, Volume III) shows the locations of the NFPD TraC surveys reported for the River Tees. For the purpose of investigating spatial variations, the River Tees has been divided into three areas: the upper, middle and lower reaches.

14.4.19 The upper reach of the River Tees is defined as the area located upstream of the Tees Barrage, whilst the middle reach falls downstream of the barrage.

² No survey took place in 2012.

The lower reaches were defined as the survey locations where sampling was undertaken at the mouth of the River Tees and the adjacent coastal waters.

Lower Reaches

- 14.4.20 The lower reaches of the River Tees have been defined as the area where Environment Agency TraC surveys were undertaken at the mouth of the river and beyond into the coastal waters (see Figure 14B-4, PEI Report, Volume III). Data for this area has been reported from 1985 to 2018, with beam trawling (2.4 m) representing the majority of surveys undertaken between 1985 and 2009. However, since 2004 seine netting has become the main fishing technique. Seine netting has the highest total catches across years, with 17,324 individuals reported; particularly high abundances were recorded in 2006, 2010, 2012 and 2014 (see Diagram 14B-2). This gear type typically captures species which utilise intertidal areas as nursery grounds and therefore, occur in high abundances (e.g. sprat, herring, and lesser sandeel).
- 14.4.21 Diagram 14B-3 shows the proportion of fish reported across all years from TraC surveys in the lower reaches of the River Tees. As expected, the majority of species recorded in this area were marine, with sprat contributing 41% to overall abundance. Herring and lesser sandeel formed the second and third largest proportions representing 15% and 10%, respectively. The following species contributed between 1% and 9% to the total proportion: plaice, dab, whiting, cod, saithe, flounder, pogge (*Agonus cataphractus*) and sandeel species (Ammodytidae). The species listed as other (a full list is included in Annex B) in Diagram 14B-3 collectively contributed 1% to the overall proportion of fish species in the lower reaches of the River Tees.
- 14.4.22 The only diadromous fish species caught in this area were European eel and sea trout. A single European eel was reported in 2004, whilst two elvers were caught in 2008. Furthermore, a total of three sea trout were recorded with a single individual recorded in each of 2004, 2009 and 2010.




Diagram 14B-2: Total fish counts for each gear type used, in Environment Agency (2019a) TraC surveys in the lower reaches of the River Tees.




Diagram 14B-3: Proportion of fish species from Environment Agency (2019a) TraC surveys in the lower reaches of the River Tees (1985 – 2018). Fish species in the legend have been presented in order of percentage contribution.

Middle Reaches

- 14.4.23 The middle reaches of the River Tees has been defined as the area upstream of the mouth of the river and downstream of the Tees Barrage (see Figure 14B-4, PEI Report, Volume III). Data from TraC surveys for this area have been reported from 1981 to 2015 (see Diagram 14B-4). Beam trawling (2.4 m) was the main gear type used from 1981 to 2004, with seine netting being the predominant method used from 2004 to 2015. High total fish counts from beam trawls (2.4 m) occurred in 1993, 1994, 2000 and 2002 where sprat, plaice and whiting dominated catches. Counts for seine netting in the middle reaches were highest between 2009 and 2011, which was predominantly due to the presence of herring and sprat in 2009 and 2010, whilst in 2011, sprat, flounder and three-spined stickleback contributed the most to the total abundance.
- 14.4.24 Sprat, plaice, whiting, and herring contributed the most to the overall abundance, representing 31%, 21%, 13% and 13%, respectively. The species contributing between 9% and 1% were: flounder, three-spined stickleback, dab, sand goby, cod, common goby, and viviparous blenny. The remaining species contributed 2% to the overall fish species proportions in the middle reaches, for a full list see Annex B.
- 14.4.25 The majority of fish species recorded in the middle reaches of the River Tees were marine; however, the diadromous species European eel, Atlantic salmon, sea trout, and river lamprey were also found to be present. Of these species, European eel was recorded most frequently with 19 individuals recorded during a total of eight sampling occasions. Five individuals of sea trout were recorded in total, with four of these being present in the past 10 years. Total counts for river lamprey and Atlantic salmon for the entire monitoring period (1981 – 2015), were three and one individuals, respectively.




Diagram 14B-4: Total fish counts for each gear type used, in Environment Agency (2019a) TraC surveys in the middle reaches of the River Tees.




Diagram 14B-5: Proportion of fish species from Environment Agency (2019a) TraC surveys in the middle reaches of the River Tees (1981 – 2015). Fish species in the legend have been presented in order of percentage contribution.

Upper Reaches

- 14.4.26 Of the fish species recorded in the upper reaches, a range of marine, freshwater, and diadromous species were reported. The upper reaches were defined as the survey area undertaken upstream of the Tees Barrage (Figure 14B-4, PEI Report, Volume III). However, the barrage was only used as an indicative reference point, as the surveys undertaken from 1982 to 1990 in the upper reaches took place prior to the construction of the barrage. All TraC surveys in the upper reaches utilised beam trawl netting (2.4 m). The highest fish counts were recorded in 1987 and 1990, with 641 and 796 fish recorded, respectively (Diagram 14B-6). In 1987 the catch was dominated by flounder and in 1990 by dace and gudgeon.
- 14.4.27 Of the fish species recorded in the upper reaches, flounder and dace contributed the most to the overall abundance of fish species, representing 60% and 20%, respectively (Diagram 14B-7). Gudgeon, European eel, roach and sprat, contributed a further 9%, 5%, 3% and 2%, respectively, whilst the remaining species combined contributed less than 1% of the total.
- 14.4.28 The diadromous fish caught were European eel and sea trout, with a total of 156 individuals of European eel caught from 1983 to 1991. Sea trout were reported in 1988 only, with one individual recorded.




Diagram 14B-6: Total fish counts for each gear type used, in Environment Agency (2019a) TraC surveys in the upper reaches of the River Tees.




Diagram 14B-7: Proportion of fish species from Environment Agency (2019) TraC surveys in the upper reaches of the River Tees (1982 – 1996). Fish species in the legend have been presented in order of percentage contribution.

14.4.29 Additional data from freshwater fish counts from NFPD Environment Agency surveys in the upper reaches of the River Tees and adjoining tributaries, recorded a total of 31,663 individuals of diadromous fish species from 1993 to 2019. Of these, a total of 24,857 individuals were counted from the River Tees and Tees Barrage. Annex C provides detailed figures showing the spatial distribution of Atlantic salmon, brown / sea trout, European eel and lamprey in the River Tees and connected tributaries.

14.4.30 Lamprey were recorded in the River Tees and River Skerne (a tributary of the River Tees) only and exhibited the smallest spatial distribution of the migratory species identified. The highest abundance of lamprey was recorded near Darlington on the River Tees but they were recorded upstream as far as Staindrop.

14.4.31 European eel had a greater distribution than lamprey, being recorded from the Tees Barrage to as far upstream as Newbiggin. European eel was also recorded on the River Lune and River Skerne, both tributaries of the River Tees. Atlantic salmon had a similar spatial extent to European eel. The highest abundances of this species were found near Eggleston, further upstream than the highest abundances of European eel and lamprey. Atlantic salmon was also recorded on the following River Tees tributaries: Egglestone Beck, Ettersgill Beck, River Lune and River Skerne. Brown / sea trout had the greatest spatial distribution of all diadromous fish species, with the highest concentrations of individuals recorded near Middleton-in-Teesdale. In addition to the tributaries where Atlantic salmon were recorded, brown / sea trout were also present in Harwood Beck and Hudeshope Beck.

14.5 Commercial Fisheries

- 14.5.1 Commercial fisheries data provides valuable information on the economic and social importance of fish stocks and whilst recognising its limitations with respect to potential bias in activities and reporting, can also be used to infer the distribution and abundance of commercial species, supplementing the fish and shellfish community baseline.
- 14.5.2 The commercial fishing activity of relevance to the study area comes from the ICES rectangle 38E8, which encompasses the fish and shellfish baseline study area for the Proposed Development. Information on commercial fishing activity in this rectangle has been collated from data reported by the MMO (2018), as part of the iFISH data system which is a UK repository where commercial fisherman (not just from the UK) are required to report administrative data under EU legislation. The data reported as part of this system gives information on the fishing activity by nationality of the vessels in the selected area, the length group of the vessel, the gear type used, the month and year of fishing and the species-specific landing weight (tonnes) and value (pound sterling).
- 14.5.3 Discussions with the NEIFCA are ongoing to identify surveillance data and any survey data held by this organisation to further supplement the commercial fisheries baseline information.

Vessel Nationalities

- 14.5.4 Table 14B-4 gives an indication of the nationality of the vessels fishing in ICES rectangle 38E8 and the landing weight recorded per annum for vessels registered in the United Kingdom (UK)³. The highest average landed weight (1,018 tonnes) was made by English vessels. Vessels from Scotland and Ireland had the second and third highest average landed weight with 85 and 39 tonnes, respectively. Welsh vessels have not reported landings from ICES rectangle 38E8 in recent years (2016 and 2017) with low annual landing weights (<10 tonnes) recorded prior to this.

Table 14B-4: Nationalities of vessels fishing in the ICES rectangle 38E8 and the weight (tonnes) of their annual landings (tonnes) (2013 – 2017).

Vessel Nationality	Landed weight (tonnes)					Average (2013 – 2017)
	2013	2014	2015	2016	2017	
UK – England	1233.7	1043.3	1082.1	1017.4	711.7	1,017.6
UK – Scotland	183.9	84.5	63.9	54.8	35.2	84.5
UK – Northern Ireland	101.9	57.7	21.6	6.5	9.1	39.3
UK – Wales	-	8.0	0.2	-	-	4.1

Source: MMO, 2018

³ Foreign nationalities are not required to report landings data via iFISH.

Fishing Activity

14.5.5 Diagram 14B-8 shows the extent of fishing activity (recorded as landed weight in tonnes) by fishing method in the ICES rectangle 38E8 for the most recent five-year period (2013 to 2017). The fishing methods identified as being used are:

- beam trawling;
- demersal otter trawling and seine netting;
- scallop dredging;
- drift and fixed netting;
- gear using hooks; and
- potting and trapping.

14.5.6 The MMO statistics showed that demersal otter trawling and seine netting were the most prevalent fishing methods operating in the ICES rectangle 38E8. From 2013 to 2017 a total of 4,369 tonnes of fish and shellfish were landed using these methods. *Nephrops* and whiting were the most targeted species, with an average landed weight of 377 tonnes and 265 tonnes, respectively. Cod, plaice, haddock and lemon sole also represented an important component of total landings by otter trawling and seine netting, representing a combined average weight of 122 tonnes.

14.5.7 The second most common fishing method used within the ICES rectangle 38E8 is potting and trapping with a total landed weight of 1,219 tonnes reported from 2013 to 2017. This method is predominately used to target lobsters and edible crabs, with an average weight of 87 tonnes and 136 tonnes landed between 2013 and 2017, respectively. In addition, but to a lesser extent, velvet swimming crab, *nephrops* and cod contributed to the total landed weight reported for potting and trapping.

14.5.8 Beam trawling, scallop dredging, drift and fixed netting, and gear using hooks, only represented a combined total of 2% of landed weight (tonnes) reported in the MMO statistics for the ICES rectangle 38E8 (2013 – 2017). Scallops comprised 88% of the total landed weight recorded for the scallop dredging fishing method, whilst mackerel dominated the reported fish catch for vessels utilising gear using hooks, representing 97% of the total landed weight. The fish and shellfish species typically targeted by drift and fixed netting were whiting and cod.

14.5.9 Diagram 14B-9 shows the annual variation in the fish and shellfish species landed weights (tonnes) from reported landings data in the ICES rectangle 38E8. In general, the total landed weights per year have remained similar, with a slight overall decline in 2017.

The species composition of the total landed weights across years was dominated by *nephrops*, with an average of 383 tonnes (2013 – 2017). This is with the exception of 2015, where whiting represented the highest total landed weight. In general, abundances were similar between years with only small variations for each species.

Nephrops, whiting and edible crabs always represented the top three landed weights.

[This page is intentionally blank]




Diagram 14B-8: Average landed weight (tonnes) (2013 – 2017) recorded in the ICES rectangle 38E8 for each gear category used. Fishing activity data taken from iFISH data system, as reported by the MMO (2018).




Diagram 14B-9: Annual landed weight (tonnes) (2013 – 2017) recorded in the ICES rectangle 38E8. Fishing activity data taken from IFISH data system, as reported by the MMO (2018).

Vessel Size

14.5.10 Table 14B-5 shows the differences in annual landed weight (tonnes) between the different vessel sizes that fish in the ICES rectangle 38E8. The vessel sizes reported are 10 m and under, and over 10 m, with similar species contributing to the overall landed weight reported from 2013 to 2017. The average landed weight reported by vessels over 10 m is dominated by *nephrops* and whiting, with an annual average of 206 tonnes and 105 tonnes, respectively. The remaining species such as edible crabs, cod and plaice also contributed to the overall landed weight, but to a lesser extent (31%). For vessels registered as 10 m and under, although the catch was still dominated by *nephrops* and whiting (177 tonnes and 163 tonnes, respectively), other species had a greater contribution to the total landed weight. For example, edible crabs, lobsters and cod contributed 15%, 12%, and 6% to the total landed weights, respectively.

14.5.11 Differences in the composition of catches landed by the two vessel sizes can be accounted for by the different gear types utilised, as shown in Diagram 14B-10. For example, the 10 m and under fleet carries out a larger amount of potting and trapping which targets edible crab and lobster. In contrast, the over 10 m fleet carries out more beam trawling and dredging, targeting whiting and scallops. However, demersal trawling and seine netting remains the dominant fishing technique for all vessels and therefore accounts for the overall dominance of *nephrops* and whiting for both sized fleets.

Table 14B-5: Length categories of vessels fishing in the ICES rectangle 38E8 and their average landed weight (tonnes) per year (2013 – 2017) for each gear type used.

Gear Type	10 m and Under	Over 10 m
Beam trawl	0.0	1.5
Demersal trawl/seine netting	475.8	398.0
Dredge	1.0	8.4
Drift and fixed netting	5.7	2.6
Gear using hooks	5.8	0.4
Pots and traps	201.1	42.7

Source: MMO, 2018

[This page is intentionally blank]




Diagram 14B-10: Average landed weight (tonnes) (2013 – 2017) recorded in the ICES rectangle 38E8 for each vessel length category. Fishing activity data taken from IFISH data system, as reported by the MMO (2018).

14.6 Species-Specific Information

Migratory Species

- 14.6.1 Diadromous species migrate between bodies of freshwater and seawater during different life phases. Major physiological changes associated with these movements occur in order to adapt to altered salinity and during such periods, sensitivity to environmental stressors increases (Shrimpton, 2012). Many migratory species, such as European eel, are also susceptible to being drawn into water intakes and outfalls as they move through estuaries and rivers (Sheridan *et al.*, 2013). For these reasons and owing to their conservation importance (see Table 14B-2), it is necessary to understand the migration patterns of the diadromous species known, or likely, to be present within the study area.

Salmon and Sea Trout

- 14.6.2 Historically, the River Tees has been subjected to serious pollution from urbanisation and industrial development, principally in the estuary and lower reaches (Moore and Potter, 2014). Since the latter part of the 20th century it has been a river in recovery, supporting a small but increasing salmon and sea trout in-river rod fishery (Environment Agency, 2009). The salmon population in the Tees has increased in the period 2004 and 2013, though numbers have become more variable. This followed a period of relatively stable abundance seen between 1995, when counts of salmon entering the Tees began, and 2003 (Moore and Potter, 2014).
- 14.6.3 Salmon are an anadromous⁴ migratory species, which during their lifetime utilises both marine and freshwater habitats. Spawning of salmon occurs in the upper reaches of rivers where females deposit eggs into nests known as 'redds' which are cut into gravelly substrate (NASCO, 2012). In the River Tees, the greatest numbers of redds are located upstream of Eggleston, with a relatively high density between Stapleton and Whorlton (Environment Agency, 2009). Once the eggs hatch, the resultant larvae remain within the interstitial gravels, utilising nutrients from the yolk sac. The larvae then develop into fry which prey on invertebrates, and then a 'parr', a young salmon distinguished by dark rounded patches evenly spaced along its sides. The length of time of the transition between life stages is geographically variable. Typically, the transition from larvae to parr occurs in the first summer in southern streams (Potter and Dare, 2003) or up to a year in upland systems. Following the parr life stage, salmon physically and morphologically change into the next life stage, known as a 'smolt' (McCormick *et al.*, 1998). This is preceding migration to the ocean following one to five years in freshwater. The migration of smolt down-river to the ocean usually occurs from spring to early summer (Thorstad *et al.*, 2012). Once salmon have spent another one to five years at sea, the adults then return to their spawning rivers, which in the UK usually peaks in June to August and October to December (Cowx and Fraser, 2003).

⁴ Migration from the sea into freshwater for spawning



- 14.6.4 Brown trout display a broad range of life history traits, including individuals that complete their lifecycle in freshwater, those that predominately inhabit estuarine waters and exhibit full anadromy (Harris *et al.*, 2017). Sea trout are therefore anadromous brown trout (*Salmo trutta*), the migratory and non-migratory forms are considered as a single species. Migration of brown trout is known to involve genetic and environmental cues, although the full trigger is not completely understood (Malcolm *et al.*, 2010). Sea trout exhibit a similar life cycle to Atlantic salmon. However, the adult marine stage of sea trout is shortened both spatially and temporally, with some immature smolt, sometimes known as 'whitling', migrating back to freshwater environments after only a very short period of time feeding at sea (usually in the first winter in the ocean), whilst 'maidens' only return to freshwater after a minimum of a year at sea (Gargan *et al.*, 2004). Adult sea trout returning to freshwater to spawn are more likely to stray from natal rivers (Degerman *et al.*, 2012; Gauld *et al.*, 2013; King *et al.*, 2016) as well as spawn over multiple years when compared to salmon. Salmonids that migrate downstream towards salt water post spawning, referred to as 'kelts', and return to freshwater to reproduce in subsequent years are likely to have increased fecundity, resulting in larger egg deposits and improved survival of offspring (Reid *et al.*, 2012).
- 14.6.5 In England and Wales there are 80 rivers which regularly support salmon, 64 of which are designated 'principal salmon rivers'. As shown in Diagram 14B-11, the performance of salmon stocks in these rivers is assessed against conservation limits (CL) which are identified by a target number of eggs deposited during spawning to ensure the status of the population remains favourable (CEFAS *et al.*, 2019).
- 14.6.6 The River Tees is included in the list of 'principal salmon rivers' and has its own 'Salmon Action Plan' (Environment Agency, 2009). The River Tees has been subject to historic pollution and is therefore recovering but supports a small rod river fishery although this has declined in recent years (Environment Agency, 2017). The River Tees is not achieving its current CL which has been identified as 14.9 million eggs. Whilst this is expected for a river in the recovery phase, it is projected that in 2021, the Tees will remain at risk of not complying with management objectives for salmon as reported by ICES (Environment Agency, 2018).
- 14.6.7 Fish monitoring surveys of salmon and sea trout, undertaken by Moore and Potter (2014), were conducted in 2008, 2009 and 2013, using acoustic telemetry to track the movements of fish. In total, 237 fish (199 salmon and 38 sea trout) were tagged with acoustic transmitters and released below the River Tees Barrage, into the lower estuary. Of these fish, 11 individuals passed through the barrage moving mainly over the barrage gates (10 fish) but also utilising the fish pass (one fish). The remaining fish either left the estuary, migrated to adjacent river systems, or were eaten by seals.




Diagram 14B-11: Main salmon rivers in England and Wales, denoting those with Salmon Action Plans (*) and those designated as Special Areas of Conservation (\$) in which salmon must be maintained or restored to favourable conservation status (CEFAS *et al.*, 2019).

14.6.8 The Environment Agency have reported monthly numbers of salmon and sea trout utilising the fish pass at the Tees Barrage since 1995; however, in 2011 the existing fish trap was changed to an electronic fish-counter to employ a non-invasive counting method (Environment Agency, 2013). It must be noted



that the counter only monitors the upstream migration of salmon and sea trout through the fish pass and therefore only represents a proportion of the run, with available alternative passage through the main barrage gates, canoe slalom, turbine fish pass and navigation lock (Environment Agency, 2019b). Furthermore, to help fish migrate, the Environment Agency and the Canal and River Trust keep the main barrage gates open as much as possible, in doing so reducing the numbers that utilise the fish passage.

- 14.6.9 The results of the monthly upstream fish count from 2011 to 2019 are presented in Diagram 14B-12. In total, 5,164 salmon and sea trout were recorded with higher numbers occurring in June to October. Typically, peak numbers of fish were recorded in either July and August and once in September (2015). Taking an average across all years between 2011 and 2019 the peak value occurred in August (mean = 207.8; std. = 238.9). Particularly high numbers of salmon and sea trout were reported in 2012 (1,661 individuals) and 2013 (1,161 individuals) however, in recent years numbers have declined with lower annual total abundances reported in 2017 (297 individuals) and 2018 (217 individuals) (2019 total annual data not complete).
- 14.6.10 Data collected from 1995 to 2011 by the Environment Agency at the Tees Barrage (presented in Diagram 14B-13 and Diagram 14B-14 (Environment Agency, 2013)), shows that mean counts of upstream migrating salmon at the Tees Barrage, peak in September and October, with mean values reported as 50.8 (std. = 54.6) and 58.6 (std. = 49.4), respectively. In comparison, peak mean sea trout counts were similar, occurring marginally later, in October, with a mean value of 119.4 (std. = 135.5).




Diagram 14B-12: Monthly combined upstream counts for salmon and sea trout from 2011 to 2019 at the Tees Barrage on the lower Tees, reported by the Environment Agency (Environment Agency, 2019b). Counter only monitors

upstream migration through the fish pass. No data was reported for January to June 2011 and August to December 2019.




Diagram 14B-13: Monthly mean upstream counts for salmon from 1995 to 2011 at the Tees Barrage on the lower Tees, reported by the Environment Agency (Environment Agency, 2013). Fish trap counts only refer to upstream migration through the fish pass. Error bars refer to standard deviation. No data was reported for January to April 1995 and July to December 2011.




Diagram 14B-14: Monthly mean upstream counts for sea trout from 1995 to 2011 at the Tees Barrage on the lower Tees, reported by the Environment Agency (Environment Agency, 2013). Fish trap counts only refer to upstream

migration through the fish pass. Error bars refer to standard deviation. No data was reported for January to April 1995 and July to December 2011.

14.6.11 The declared rod catches for salmon and sea trout in the River Tees from 2007 to 2017 are presented in Diagram 14B-15, showing that for both species there has been a general decline in overall rod catches in recent years. This is likely due to actions taken by the Environment Agency and Defra to reduce the overall exploitation of salmon and sea trout in England and Wales, such as catch and release measures (such as: restrictions on night fishing; size limits, for trout only; and bag limits, for trout only) (Environment Agency, 2017). Salmon rod catches in this period were highest in 2008 with 267 fish, which declined to 2014 when only 16 fish were declared. There has been a general increase since 2014 to 2017, when 67 salmon were caught. Sea trout in contrast, peaked in 2014, with 114 fish catches declared. Total sea trout rod catches then declined in 2015 where there was a three year low in total catches, ranging from 11 to 13 fish. It must be noted that data from the rod and line fishery does not necessarily reflect abundance as it is not corrected for other variables, such as effort. In addition, some migration occurs outside of the angling season. However, this data is useful for a general indication of historical trends.




Diagram 14B-15: Summary of salmon and sea trout rod catches from the river Tees that were declared from 2007 to 2017, as reported by the Environment Agency (2017).

14.6.12 Only one individual of salmon was recorded in Environment Agency TraC surveys in the River Tees, being captured in 1998 in the middle reaches of the river. Nine individual sea trout were recorded in the TraC surveys, being recorded mainly in the last 10 years (six individuals) in both the lower and middle reaches of the river. One individual was recorded in the upper reaches in 1998. On average sea trout individuals measured 320 mm ($n = 8$), with the smallest measuring 87 mm (2011), whilst the largest measured 635 mm (2004).

European Eel

- 14.6.13 European eel is a catadromous migratory species, whose spawning occurs in the Sargasso Sea where the adults subsequently die. The newly hatched larvae, known as leptocephali, are transported to the continental shelf of the North Atlantic by the prevailing currents of the Gulf Stream, where they metamorphose into the life stage of glass eel and subsequently, in freshwater and coastal waters become pigmented 'elvers' (Aerstrup *et al.*, 2009; Potter and Dare, 2003). Eels migrate upstream into freshwater predominately during spring but may continue to do so until early Autumn (ICES, 2010). Once within freshwater habitats, eels remain for five to 15 years, transforming into yellow eels and then finally to silver eels when they begin their downstream migration through rivers and estuaries towards spawning grounds, predominately between August and December (Behrmann-Godel and Eckmann, 2003; Tesch, 2003; Chadwick *et al.*, 2007). Spawning occurs mainly in spring (Righton *et al.*, 2016). However, it must be noted that some eels do not migrate into freshwater but instead inhabit estuaries as 'elvers' and yellow eels before returning to spawning grounds.
- 14.6.14 Throughout England, European eels are present in almost all rivers, although in recent years their numbers have dramatically declined. This has resulted in European eels being listed as 'critically endangered' on the IUCN Red List since 2008. There are multiple reasons for the decline of European eel numbers, including barriers to migration, hydropower turbines, loss of wetland, and the introduction of the parasitic nematode *Anguillicola crassus* (UK BAP, 2012). The River Tees Barrage has the potential to act as a barrier but has built opportunities for the migration of glass eels into its design, although the escapement of adult silver eels around the barrage is unknown.
- 14.6.15 The current population size and distribution of European eels in the River Tees is unknown. However, European eel have been reported in Environment Agency TraC surveys in the River Tees (see Diagram 14B-16), with a total of 178 individuals found in total across all surveys. The majority of European eel were recorded in the upper reaches of the River Tees. Only three individuals of European eel were recorded in TraC surveys in the River Tees, exhibiting a total length of 280 mm (2008), 282 mm (2008), and 625 mm (2004). These fish were all recorded in the lower reaches of the River Tees.




Diagram 14B-16: Total fish counts for European eel (*Anguilla anguilla*), in Environment Agency (2019a) TraC surveys in the three reaches of the River Tees.

Sea and River Lamprey

- 14.6.16 Sea lamprey and river lamprey are both anadromous migratory species. Both species spawn in spring and early summer in freshwater, followed by the ammocoetes life stage (larval phase) which is spent in the silt beds of streams and rivers (Laughton and Burns, 2003). Following spawning, all adult individuals die (Maitland, 2003). In the ammocoetes phase, lamprey feed on organic detritus and can spend several years in this life stage, eventually transforming into the adult life stage in late summer and onwards (Laughton and Burns, 2003). Once an adult, both river and sea lamprey migrate out to sea where they become parasitic, using their suckers to attach onto host fish and feed on their blood (Maitland, 2003). The adults then return to freshwater once they have spent several years in the marine environment (Laughton and Burns, 2003).
- 14.6.17 Sea lamprey is widely dispersed at sea as they are solitary feeders, being rarely found in coastal and estuarine waters (Moore *et al.*, 2003). The distribution of sea lamprey is chiefly defined by their host (Waldman *et al.*, 2008) and they are often found at considerable depths in deeper offshore waters (Moore *et al.*, 2003). Sea lamprey typically feed on the blood of a range of marine mammals and fish, which include, herring, cod, pollack, Atlantic salmon, haddock, shad, and basking sharks (Kelly and King 2001, ter Hofstede *et al.*, 2008). In contrast, river lampreys are usually found in coastal water, estuaries and accessible rivers and young river lamprey are often found in large congregations (Maitland *et al.*, 2003). River lamprey feed on a variety of estuarine fish, predominantly herring, sprat and flounder. River lamprey generally spend one to two years in estuaries and in the



autumn, between October and December, stop feeding and move upstream (Natural England, 2010). Sea lamprey normally migrate into freshwater in April and May as adults, whilst the migration to sea can vary from river to river, although the metamorphosis of larvae into adults, occurs in July and September (Maitland, 2003).

- 14.6.18 The UK distribution of river lamprey and sea lamprey, presented in Diagram 14B-17, shows that both species have been recorded in the River Tees. River lamprey were also recorded in Environment Agency TraC surveys of the River Tees, with three individuals being recorded in the middle reaches in 1992. Although Diagram 14B-17 suggests sea lamprey are present in the River Tees, this species has not been recorded during the Environment Agency TraC surveys to date.




Diagram 14B-17: UK Distribution of River Lamprey (left) and Sea Lamprey (right) (JNCC, 2013a and 2013b).

Pelagic Fish Species

Herring

- 14.6.19 Herring is an important commercial species and represents a significant prey species for many predators, including large gadoids (such as cod), dogfish, sharks, marine mammals and birds (ICES, 2006a). Herring is a pelagic fish and is found mostly in continental shelf areas to depths of 200 m (Whitehead, 1986). Juveniles are generally distributed separately from adults, being found in shallower water, migrating into deeper waters to join the adult stock after two years. In the North Sea 1-group herring are restricted within the 100 m depth contour and are most abundant in the south east, Kattegat and along the British east coast (ICES, 2006a).

- 14.6.20 Herring exhibits multiple geographically distinct stocks in UK waters (Tappin *et al.*, 2011), with three major populations which are identified with different spawning times. The major population associated with the study area is the Banks population which is located in the Central North Sea and off the English coast, with spawning occurring from August to October (Ellis *et al.*, 2012). Herring are demersal spawners, which means when spawning occurs, large numbers of eggs are released (~50,000 per female) near the seafloor, which sink and attach to gravel, stones and shell where they form a dense mat. Spawning can occur in episodes which are weeks apart (Dempsey and Bamber, 1983). Once developed into juvenile fish, herring aggregate into shoals which migrate into estuaries and shallow waters where they remain for six months to a year (Dipper, 2001). The study area is recognised by Ellis *et al.* (2012) as being a high intensity nursery ground for herring.
- 14.6.21 Herring larvae usually consume copepods and other small planktonic organisms (Daan *et al.*, 1985), whilst juveniles will predominantly feed on calanoid copepods (Blaxter and Hunter, 1982) as well as euphausiids, hyperiid amphipods, juvenile sandeels, *Oikopleura* spp., and fish eggs (Last, 1989). Larger adult herring consume predominantly copepods as well as small fish, arrow worms, and ctenophores.
- 14.6.22 The herring fish stock in the North Sea is caught mainly using beam or pelagic trawls (Cotter *et al.*, 2004) and is used for human consumption or constitutes bycatch from industrial fisheries. In the North Sea, Skagerrak and Kattegat and eastern English Channel, ICES landings data for herring reported 602,328 tonnes in 2018. Herring did not represent one of the main fish species targeted in the ICES rectangle 38E8, with an average landed weight of only 1.2 tonnes for the period 2013 to 2017 (MMO, 2018). Herring were chiefly caught using the method of demersal trawling and seine netting from vessels of both size categories (10 m and under and over 10 m).
- 14.6.23 In Environment Agency TraC surveys in the River Tees, herring was recorded in all reaches of the River Tees, although only one individual was recorded in the upper reaches (caught in 1986). Herring were predominantly found in the past 10 years (largely as a consequence of changes to the monitoring programme) and represented the second highest proportions of fish found in the lower reaches and the fourth highest in the middle reaches, contributing 15% and 13% to the total fish composition, respectively.
- 14.6.24 The population structure of herring in the North Sea, usually comprises two distinct groups from 90 – 190 mm and 190 – 300 mm (ICES, 2006a). The lengths recorded for herring as part of the TraC surveys in the River Tees are shown in Diagram 14B-18, which demonstrated a distinct group of individuals from 55 – 65 mm, representing 1-group fish, which is typical in coastal areas and confirms use of the area as a nursery ground. The overall mean length recorded was 67 mm (\pm std. = 14 mm), with the majority of individuals measuring between 45 mm and 95 mm. The smallest recorded length was 36 mm with one individual recorded in 2013 and one individual recorded in 2004. The largest size capture was an individual measured at 218 mm, which was caught in 2002.




Diagram 14B-18: Size classes (mm) (based on total length (mm)) measured for herring (*Clupea harengus*) in Environment Agency (2019a) TraC surveys in the River Tees. Data has been presented for the first 95% of fish counts.

Sprat

- 14.6.25 Sprat is a short-lived, small-bodied pelagic schooling species that is relatively abundant in shallow waters. Sprat are an important food resource for a number of commercially important predatory fish, seabirds and marine mammals. Sprat themselves primarily feed on copepods, cladocerans, sea-squirts, bivalve larvae, mysids and euphausiids (Maes and Ollevier 2002; ICES, 2005).
- 14.6.26 Spawning of sprat is thought to occur from May to August, peaking in May to June (Coull *et al.*, 1998), with batches of eggs released repeatedly throughout the spawning period (Milligan, 1986). Spawning occurs in coastal waters up to 100 km offshore, and in deep basins (Whitehead, 1986; Nissling *et al.*, 2003). Once released, the eggs and larvae, which are pelagic, move into coastal nursery areas by larval drift (Hinrichsen *et al.*, 2005; Nissling *et al.*, 2003). The study area is recognised by Coull *et al.* (1998) as being an important nursery ground for sprat, with spawning grounds located nearby.
- 14.6.27 Sprat is particularly abundant in shallow areas in the North Sea and is fished commercially, primarily for use in fish meal and as bait (ICES, 2006b). In the central North Sea (ICES Subarea 4b) official landings of sprat in 2018 totalled 179,664 tonnes (ICES, 2019a). However, in the ICES rectangle 38E8, sprat is not an important commercial species, with no reported landed weights of sprat for this area (MMO, 2018).
- 14.6.28 In Environment Agency TraC surveys in the River Tees, sprat was recorded in all reaches of the river, with the highest relative abundances being recorded in the lower reaches with 10,593 individuals across all years

(1981 – 2018), with a further 7,096 individuals recorded in the middle reaches between 1981 and 2015. In the upper reaches, sprat was recorded in only three years: 1986, 1987, and 1996. In the lower and middle reaches, sprat was recorded as present in all years sampled, excluding 2002, and represented the highest proportions of fish species found, contributing 41% and 31% to the total proportion of fish, respectively.

14.6.29 In the North Sea, landings of sprat are dominated by 0-group, 1-group, and 2-group fish, with lengths ranging from 50 – 100 mm (ICES, 2006b). The lengths recorded for sprat as part of the TraC surveys in the River Tees are shown in Diagram 14B-19, with the mean length recorded as 61 mm (\pm std. = 31 mm), with most individuals measuring between 40 mm and 80 mm. The smallest recorded length was 31 mm (recorded in 2010), whilst the largest size recorded was 136 mm, for which one individual was caught in 2015.




Diagram 14B-19: Size classes (mm) (based on total length (mm)) measured for sprat (*Sprattus sprattus*) in Environment Agency (2019a) TraC surveys in the River Tees. Data has been presented for the first 95% of fish counts.

Mackerel

14.6.30 Mackerel is a widely distributed migratory fish in the North Atlantic and is one of the most abundant fish species in this area (ICES, 2011). Mackerel spend their entire life in the pelagic environment and are an important food source for sharks, tuna and dolphins (Tappin *et al.*, 2011). This species is also exploited by commercial fisheries, which in the past has caused the collapse of abundant stocks in the North Sea (ICES, 2006c).

14.6.31 Mackerel in the eastern Atlantic is divided into three spawning components, the North Sea being one of these (ICES, 1999). Mackerel overwinter in deeper waters, which in the North Sea typically occurs in Skagerrak and the north-eastern North Sea (Jansen and Gislason, 2011). They then move



closer to the shore in spring when temperatures range between 11°C and 14°C. The main spawning period for mackerel occurs in mid-May to late June, taking place particularly in the central North Sea (Jansen and Gislason, 2011). After this period, mackerel redistribute in the North Sea or migrate into surrounding waters. Mackerel are batch spawners (Murua and Saborido-Rey, 2003), reported as having a fecundity of between 130,00 and 1,100,000 eggs in the North East Atlantic (Macer, 1976).

- 14.6.32 Mackerel feed on zooplankton, in particular large quantities of pelagic crustaceans, and the pelagic larvae and juvenile stages of numerous commercially important fish species (ICES, 2008) which includes smaller fish such as sprat, herring and sandeel (Wheeler, 1978). During the winter, mackerel spend most of their time on the seabed where they are not active feeders.
- 14.6.33 Mackerel were identified as being an important commercial species in the ICES rectangle 38E8 (see Section 14.5), with average recorded landed weights in this area equating to 20 tonnes per annum (from 2013 – 2017) (MMO, 2018). They are also of commercial importance for the North Sea (ICES Subarea 4) as a whole, although official landings data of mackerel in the North Sea was unavailable (ICES, 2019b). Only one individual was recorded in the River Tees in Environment Agency TraC surveys, occurring in the lower reaches of the river in 2006 (see NFPD Fish Counts – Environment Agency in section 14.4)).
- 14.6.34 Mackerel can reach a maximum size of 65 cm, although are rarely found larger than 40 cm (Tappin *et al.*, 2011). The one individual recorded in the River Tees was an adult, measuring 386 mm in total length.

Demersal Fish Species

Cod

- 14.6.35 Cod is widely distributed throughout the North Sea (Heessen, 1993) and is found in shallow coastal waters to the shelf edge (200 m depth). Juveniles prefer shallow waters where there are a range of habitats such as seagrass beds, gravel, rock, and boulders which provide protection from predators. Cod are omnivorous, but in the North Sea are predominantly piscivorous, feeding on young clupeids, flatfish and gadoids (Daan, 1973). The typical feeding times for cod are dawn and dusk (Cohen *et al.*, 1990).
- 14.6.36 From late winter to early spring, adult cod migrate to offshore spawning grounds, typically at depths of 20 m to 100 m in the North Sea (often 200 m elsewhere) (Dipper, 2001). Cod is classified as a determinate multiple spawner (McEvoy and McEvoy, 1992) with experiments reporting between eight and 22 batches spawned per season (Kjesbu *et al.*, 1992). In a single spawning event, females produce between three million and six million eggs (Trippel, 1998), which rise to the surface and drift with ocean currents (Dipper, 2001).
- 14.6.37 The eggs and larvae of cod remain in the water column, developing into juvenile fish within six months. When individuals reach a size of approximately 7 cm, juveniles move to the sea bed where they become



demersal, often occurring between July and August (Heessen and Daan, 1994). Juvenile cod then move into coastal nursery areas once the spawning season is over, with young cod often found in estuaries and shallow waters. The study area is recognised by Ellis *et al.* (2012) as being a high intensity nursery ground for cod.

14.6.38 Cod was identified as being an important commercial species in the ICES rectangle 38E8 (see Section 14.5), where on average, 57 tonnes of cod were recorded per year as landed from 2013 to 2017, representing 5% of the total landed weight for all species and years in the area (MMO, 2018). In this area, cod was caught mainly from vessels that were 10 m and under in size, using trawling and seine netting. Overall in the North Sea (ICES Subarea 4), ICES official landings for cod in 2018 were 35,789 tonnes (ICES, 2019c). In Environment Agency TraC surveys in the River Tees, cod was recorded in both the middle and lower reaches of the River Tees but not in the upper reaches. Cod was recorded as present in most years and represented 3% and 2% of the total proportion of fish species found in the lower and middle reaches, respectively.

14.6.39 In the North Sea, the population structure of cod is dominated by individuals aged one (100 – 250 mm) and two years (200 – 400 mm) old (ICES, 2006d). The lengths recorded for cod as part of the TraC surveys in the River Tees are shown in Diagram 14B-20, with the mean length recorded as 151 mm (\pm std. = 48 mm) with the majority of fish being one year of age. The smallest recorded length was 30 mm. The largest size recorded was 504 mm, which was caught in 2009.




Diagram 14B-20: Size classes (mm) (based on total length (mm)) measured for cod (*Gadus morhua*) in Environment Agency (2019a) TraC surveys in the River Tees. Data has been presented for the first 95% of fish counts.

Whiting

- 14.6.40 Whiting is most abundant in water depths of 30 m to 100 m, but is very common in shallow waters (Wheeler, 1978). Whiting are a benthic-pelagic species, found in association with a variety of substrates including mud, gravel, sandy and rocky areas, (Barnes, 2008). The spatial distribution of whiting, particularly in the northern North Sea, appears to be affected by sea surface temperature (Zheng *et al.*, 2002).
- 14.6.41 Spawning of whiting is mainly triggered by temperature (5 – 10°C is optimal) and takes place from February to June (Coull *et al.*, 1998), peaking in spring in shallow waters (Wheeler, 1978). The study area is recognised by Ellis *et al.* (2012) as being a high intensity nursery ground for whiting. Once spawned, whiting grows relatively slower during their first year of life, with large variations in growth rates between individuals (Hislop *et al.*, 1991). Whiting typically reach maturity after two years and often spawn during this year.
- 14.6.42 A female whiting of 30 cm will produce 400,000 eggs, which compared to other gadoids, demonstrates high relative fecundity (Hislop and Hall, 1974). Whiting release their eggs in many batches over a period which usually lasts up to 14 weeks (Hislop *et al.*, 1991).
- 14.6.43 Whiting is a common predator of other fish species, feeding predominantly on commercially important species, including its own offspring. When greater than 30 cm, whiting feed almost entirely on fish, including younger age classes of herring, cod and haddock as well as smaller species such as sprat, sandeel, and Norway pout (Hislop *et al.*, 1991). When in the pelagic larvae life stage, whiting feed on the nauplii and copepodite stages of copepods (Last, 1978) whilst juvenile whiting (< 20 cm) feed on crustaceans, including euphausiids, mysids and crangonid shrimps.
- 14.6.44 In the North Sea area, the population structure of whiting is typically dominated by high relative abundances of those aged 1 and 2+ years. This same pattern was recorded in TraC surveys in the River Tees, with two peaks in length frequency illustrated in Diagram 14B-21. The two clear size groups comprise a peak from 105 mm to 145 mm and a second but smaller peak at 250 mm to 265 mm. The smallest recorded length for whiting across all surveys was 42 mm, caught in 2012, whilst the largest length was 382 mm, which was recorded in 2008. The mean total length for whiting was 186 mm (\pm std. = 83 mm).




Diagram 14B-21: Size classes (mm) (based on total length (mm)) measured for whiting (*Merlangus merlangus*) in Environment Agency (2019a) TraC surveys in the River Tees. Data has been presented for the first 95% of fish counts.

Dover Sole

- 14.6.45 Dover sole is a southern species whose northern limit is in the North Sea. Sole favours sandy and sandy muddy substrates, which they can bury into, in waters of up to 50 m depth. The spatial distribution of Dover sole varies between life stages, with juveniles favouring coastal nursery grounds whilst older and larger individuals occupying deeper offshore waters (Teal, 2011).
- 14.6.46 Spawning in the North Sea typically occurs in March to June and peaks in April (Tappin *et al.*, 2011). Spawning takes place inshore, generally in estuaries, where high numbers of eggs are released (up to 500,000 eggs). The eggs then drift into high productivity shallow sandy nursery grounds which provide a good feeding ground for juveniles (Dipper, 2001). Dover sole will inhabit nursery grounds for up to two years before migrating offshore (Rijnsdorp *et al.*, 1992). The study area is not recognised by Ellis *et al.* (2012) or Coull *et al.* (1998) as being a Dover sole nursery ground or spawning area.
- 14.6.47 Dover sole is a nocturnal feeder spending the day buried in bottom sediment (Kruuk, 1963). Dover sole are olfactorial feeders with multiple sensory organs located on their blind side which aid prey detection. Juveniles and adults typically eat polychaete worms and small echinoderms and small molluscs (Braber and Groot, 1973), whilst pelagic larvae feed on copepod nauplii (Russel, 1976).
- 14.6.48 Sole is mainly found in the southern and eastern North Sea, south of the line from Flamborough (ICES, 2006e). In summer and autumn this line represents a steep gradient in temperature which divides the North Sea into



a cold stratified northern section and a warm mixed southern section where the bottom waters are also warmer (ICES, 1965). Official landings of Dover sole in the North Sea (ICES Subarea 4) for 2018 estimated by ICES was 11,199 tonnes. In the ICES rectangle 38E8, Dover sole represents only a small proportion of the total landed weight from 2013 to 2018, that being 0.8% with an average landed weight (tonnes) per year of 9.5 tonnes (MMO, 2018). Fishing activity for Dover sole in this ICES rectangle uses predominantly demersal trawls and seine netting by vessels which were typically 10 m and under in length.

- 14.6.49 In Environment Agency TraC surveys in the River Tees, Dover sole was recorded occasionally in the lower reaches of the river and to a lesser extent in the middle reaches but was never caught in the upper reaches. A total of 19 fish were recorded in these surveys from 1991 to 2016.
- 14.6.50 In the North Sea, Dover sole that are over 10 years of age are rarely caught, with length distributions dominated by individuals that are one year old (~100 mm) and a second peak at two years old (~200 - 250 mm) (ICES, 2006e). In TraC surveys in the River Tees, the majority of individuals were two years old with the mean total length recorded being 251 mm (\pm std. = 97 mm). The smallest recorded length of sole was 44 mm (caught in 2013) whilst the largest length was 410 mm (caught in 2002).

Plaice

- 14.6.51 Plaice are found on all UK coasts, normally located on sandy substrata, as well as gravel and mud, and are frequently observed on sand patches in rocky areas (Tappin *et al.*, 2011).
- 14.6.52 Female plaice usually mature at four to five years old, whereas males typically become sexually mature at two to three years of ages (Heessen and Rijnsdorp, 1989). Plaice generally spawn between January and April, at depths of between 20 m and 40 m, with up to 500,000 eggs being released. Spawning grounds in the North Sea with high concentrations of egg production include Southern Bight and eastern Channel, whilst further north on the east coast of the UK, spawning is less intense (Ellis *et al.*, 2012). Coastal and inshore waters of the North Sea however, represent important nursery areas (Kuipers, 1977) although the study area is recognised by Ellis *et al.* (2012) as being only a low intensity nursery ground for plaice. Following spawning, plaice reach their peak densities in May, and in June and July older fish tend to migrate offshore, whilst juveniles remain in the intertidal zone until autumn (Kuipers, 1997). The planktonic larvae stage lasts four to six weeks, following which plaice settle on sandy substrates within nursery grounds.
- 14.6.53 Plaice usually feed during the day although during spawning periods adults do not feed at all (De Clerk and Buseyne, 1989). Adult plaice will often eat polychaete worms and bivalve molluscs, but also feed on small crustaceans, such as amphipods, mysids, and small shrimp (De Clerk and Buseyne, 1989). The diet of plaice larvae usually consists of pelagic tunicates although copepods, algae and bivalve post-larvae are also consumed (Ryland, 1964).

14.6.54 Plaice is an important commercial species in the North Sea, with estimated landings of plaice of 50,783 tonnes being reported by ICES (ICES, 2019d). In the ICES rectangle 38E8, plaice represented 3% of the total landed weight (tonnes) of all fish species caught from 2013 to 2017, with the main fishing technique used being demersal trawling and seine netting (MMO, 2018). In Environment Agency TraC surveys in the River Tees, plaice was recorded in both the middle and lower reaches of the River Tees but not in the upper reaches. The highest relative abundances were recorded in the middle reaches, with 4,741 fish caught across all surveys (compared to 2,248 individuals in the lower reaches). Plaice was recorded as present in most years and represented 9% and 21% of the total proportion of fish species found in the lower and middle reaches, respectively.

14.6.55 In the North Sea, the typical length frequency distribution is dominated by two peaks, corresponding with age groups 1 (~100 mm) and 2 (~200 – 250 mm), the latter being the more dominant (ICES, 2006f). In the River Tees, the lengths frequency distribution showed only one peak from 40 – 100 mm, the mean length being 79 mm (\pm std. = 39 mm) (Diagram 14B-22). The smallest recorded length was 13 mm, recorded in 2014 whilst the largest size recorded size was 418 mm, which was caught in 2007.




Diagram 14B-22: Size classes (mm) (based on total length (mm)) measured for plaice (*Pleuronectes platessa*) in Environment Agency (2019a) TraC surveys in the River Tees. Data has been presented for the first 95% of fish counts.

Sandeel

14.6.56 Five sandeel species occur in the North Sea, including Raitt's sandeel (*Ammodytes marinus*) which is the most common although the lesser sandeel (*Ammodytes tobianus*) and great sandeel (*Hyperoplus lanceolatus*) are also prevalent. Sandeel are an important element of the food chain in the North Atlantic and feed other fish species, sea birds and marine mammals,



including cod, herring, salmon, mackerel, grey seal, harbour seal, porpoise and puffins (Dipper, 2001).

- 14.6.57 Sandeel spend a large proportion of the year buried in the sediment, only emerging into the water column to spawn briefly in winter (November to February), and for an extended feeding period during the spring and summer months (van der Kooij *et al.*, 2008).
- 14.6.58 The distribution of sandeel (referring to all species within the genus *Ammodytes* spp) is highly patchy due to their preference for sandy habitats, favouring coarse sand with fine to medium gravel and a low silt content (Holland *et al.*, 2005; Greenstreet *et al.*, 2010). Populations are also associated with seabed morphological features such as subtidal sandbanks (MarineSpace *et al.*, 2013). However, this species is broadly found from inshore waters down to the shallow sublittoral zone (i.e. to 70 m depths).
- 14.6.59 During spawning, females will release between 4,000 and 20,000 eggs (Tappin *et al.*, 2011), which they deposit within the sandy habitat of the adults as they are demersal spawners (Winslade, 1971). Once hatched, the larvae are pelagic, spending their time in the water column until they develop into juveniles in the winter when they burrow into the sediment (Macer, 1966). The study area was not recognised by Ellis *et al.* (2012) as being within a nursery ground for sandeel.
- 14.6.60 Sandeel, which is mainly used for fish meal, supports one of the largest fisheries in this area. In the northern and central North Sea (ICES Divisions 4.a and 4.b) sandeel fisheries have been divided into 'Sandeel Areas', Sandeel Area 4, in the northern and central North Sea (ICES divisions 4.a and 4.b), being associated with the study area (ICES, 2019e). Official landings of sandeel in Sandeel Area 4 for 2018 estimated by ICES was 42,526 tonnes. No landings of sandeel from ICES rectangle 38E8 has been reported for 2013 – 2017 (MMO, 2018).
- 14.6.61 In Environment Agency TraC surveys in the River Tees, no sandeel were recorded in the upper reaches of the River Tees and only six individuals were present in the middle reaches. The majority of sandeel were found in the lower reaches of the river, being caught using seine netting. Lesser sandeel were the *Ammodytidae* species caught in the highest abundances, whilst Corbin's (*Hyperoplus immaculatus*) and greater sandeel were also recorded. In the past 10 years, the total relative abundance of sandeel was 2,056 individuals. Lesser sandeel in the lower reaches of the River Tees, composed 10% of the total proportion of fish reported from the TraC surveys.
- 14.6.62 The total lengths of sandeel recorded in TraC surveys in the River Tees is illustrated in Diagram 14B-23. The length at maturity for lesser sandeel ranges from 110 - 150 mm. Thus, with mean total length for sandeel of 98 mm (\pm std. = 18 mm), the majority of fish measured were immature. The smallest recorded length for sandeel across all surveys was 64 mm, caught in 2010, whilst the largest length was 260 mm, which was a lesser sandeel recorded in 2005.
- 14.6.63 In December 2019, a benthic grab survey was carried out within the study area (see Appendix 14D: Subtidal Benthic Ecology). From these samples, a



total of 11 sandeels (*Ammodytes* spp) were recorded across six sites located within the study area. Individuals ranged from 31 mm to 197 mm in length representing both juveniles and adults. The presence of juveniles suggests that the study area is used by sandeels as a nursery ground with possible spawning occurring within the area.




Diagram 14B-23: Size classes (mm) (based on total length (mm)) measured for *Ammodytidae* (sandeel species) in Environment Agency (2019a) TraC surveys in the River Tees. Data has been presented for the first 95% of fish counts.

Shellfish

Nephrops

14.6.64 *Nephrops* (Norway lobster) are distributed according to the extent of cohesive muddy sediments, in which they construct their burrows (Howard, 1989). The type of sediment also dictates the structure of the *Nephrops* populations, with areas of sandy mud having higher population densities and smaller sized individuals, whilst fine sediment is characterised by lower densities of larger-bodied individuals. Males can reach up to sizes of 250 mm in length and mature at three years of age (Tappin *et al.*, 2011). Mating of *Nephrops* in the UK occurs in summer after the female has moulted, the eggs being laid on their pleopods in the autumn (Farmer, 1974). The eggs stay on the pleopods for up to 60 days in temperate waters, releasing a pelagic larva, after which the juveniles metamorphose into burrowing post-larvae.

14.6.65 *Nephrops* in the North Sea is targeted mainly by demersal trawlers but are also caught using seine nets and pots (Cotter *et al.*, 2004). In the North Sea, the *Nephrops* fishery has been divided into different functional units by ICES, the unit relating to the study area being Farn Deep in the central North Sea (ICES Division 4.b, Functional Unit 6) (ICES, 2019f). Since April 2016, a

range of measures on UK vessels fishing for Norway lobster in Farn Deep were implemented by the UK in order to reduce fishing mortality on the stock and unsustainable fishing of the stock. The official total landings weight of *Nephrops norvegicus* in Farn Deep was 1,807 tonnes in 2018.

- 14.6.66 *Nephrops* represents an important commercial species in the ICES rectangle 38E8, having the highest total landed weight of all fish and shellfish for the area, representing 34% of the total from 2013 to 2017 (MMO, 2018). An average landed weight of 383 tonnes per annum (2013 – 2017) was recorded for *Nephrops* with demersal trawling and seine netting being the main gear types used to catch this species (representing 98% of the total landed weight). However, pots and traps also contributed (1%) to the overall fishing activity of *Nephrops*, with an average landed weight of 5 tonnes per annum. Landings of *Nephrops* were from vessels that were both 10 m and under and over 10 m in size, those representing over 10 m contributing 54% to the average landed weight (2013 – 2017).

Edible crab

- 14.6.67 Edible crab in the North Sea is found in waters between 25 m and 300 m, with a preference for bedrock, mixed coarse grounds, and offshore in muddy sands (Neal and Wilson, 2008). Edible crabs copulate in the spring and summer, the female crabs becoming gravid, carrying their eggs under the abdomen. At this time, females do not feed and remain in pits in the sediment or under the rocks and therefore fishing pressure is unlikely to affect the larval population (Howard, 1982). In the North Sea, brooding females migrate offshore to release the larvae, which once hatched remain in the water column for between 60 and 90 days before settling. Tagging surveys off the coast of Norfolk, have shown that mature females undertake long-distance northerly migrations to the Yorkshire coast, although more recent studies suggested this may be a discrete population of edible crabs (Eaton *et al.*, 2003).
- 14.6.68 On the coast of England and Wales, edible crabs represent a significant commercial fishery, with the most recent reported landings in 2015 above 14,000 tonnes (ICES, 2016). The fishery in England and Wales is mainly targeted by vessels under 12 m, which utilise pots and operate close to the coast. Average numbers of edible crabs reported from CEFAS (and ICES in parenthesis) samples in the central North Sea from 2014 to 2016 was 3,703 (2,440) (n = 152 (77)) (CEFAS *et al.*, 2017a).
- 14.6.69 Traditional fisheries of edible crabs are located off Yorkshire and Northumberland and are an important fishery in the ICES rectangle 38E8 (MMO, 2018). From 2013 to 2017, the annual average recorded landed weight of crab in this area was 269 tonnes, which represents the third highest average landed weight by species in that area (12% of the total landed weight in the rectangle). The annual reported landed weights (2013 – 2017) of edible crab in the ICES rectangle 38E8 have remained similar between years, peaking in 2014 at 135 tonnes. Pots and traps were the main gear types used to catch edible crab, with 99% of the average landed weight being recorded using these methods. Landings of edible crab were also mostly from vessels that were 10 m and under in size, representing 75% of the average landed weight.

Lobsters

- 14.6.70 European lobster is generally found from the intertidal zone to depths of 60 m. This species exhibits site fidelity although home extents can range between 2 km and 10 km (Bannister *et al.*, 1994). Lobsters are solitary animals and inhabit holes and tunnels that they build below rocks and boulders (Wilson, 2008). Females can spawn annually or following a bi-annual pattern, with reproduction taking place during the summer (Atema, 1986). They do not make extensive migrations when berried and hatching takes place in spring and early summer on the same grounds (Pawson, 1995).
- 14.6.71 Similar to edible crab, European lobster is an important commercial shellfish species caught along the coasts of England and Wales, with the most recent annual landings reported in 2015 being 1,885 tonnes (ICES, 2016). The gear types used to target lobster include pots and traps, predominantly by vessels under 12 m in size. The most productive lobster fishery area is Yorkshire and Humber, where landings represent 44% of the total in 2015, and have increased progressively since 2006 despite fishing effort remaining similar since 2011. Lobster fisheries in England have been divided into six Lobster Fisheries Units (LFU), the study area falling within the Yorkshire Humber LFU but also associated with the Northumberland LFU (CEFAS, 2017b). Both LFUs have high levels of exploitation which are above the maximum reference point limit, although this has decreased in recent years (CEFAS, 2017b). Furthermore, fishing pressure is particularly high around the Minimum Landing Size (CEFAS, 2017b).
- 14.6.72 From 2013 to 2017, the average annual recorded landed weight of lobster in the ICES rectangle 38E8 represented 8% of the total landed weight across the years and equated to 88 tonnes (MMO, 2018). Pots and traps were the main gear types used to catch lobster, with 99% of the average landed weight being recorded using these methods. Landings of lobster were also mainly from vessels that were 10 m and under in size, representing 91% of the average landed weight.

14.7 Baseline Evolution

- 14.7.1 Baseline conditions for fisheries and fish ecology can be influenced by a variety of factors including, commercial exploitation, pollution, loss of habitats and food resources due to riverine and coastal development, conservation and management measures, and global warming (i.e. climate change). These factors can not only influence the abundance and distribution of species but also life history processes including growth and reproduction.
- 14.7.2 Monitoring of fish communities within the Tees Estuary by the Environment Agency has been ongoing since the 1980s (Environment Agency, 2019a; 2019b). However, owing to changes in sampling methods, locations and sampling frequency during the monitoring period, it is not possible to reliably identify any long-term trends to inform this assessment of baseline evolution.
- 14.7.3 Within the study area, climate change impacts due to factors such as increasing sea surface levels and warming sea temperatures is considered



to be one of the principle ways in which baseline conditions are likely to evolve during the life cycle of the Proposed Development and is therefore considered in further detail below.

- 14.7.4 Future UK Climate Projections 2018 (UKCP18) from the Met Office for the Stockton-on-Tees area (The Met Office, 2019), based on a 1981 – 2000 baseline⁵, uses a range of possible scenarios, classified as Representative Concentration Pathways (RCPs), to inform different future emission trends. RCP 8.5 has been used for the purposes of this assessment as a worst-case scenario.
- 14.7.5 Based on RCP 8.5, there is a 50% probability that sea levels will have risen 8 cm by 2022 (commencement of construction) and 11 cm by 2026 (commencement of operation). By 2051 (i.e. the end of the Proposed Developments operational lifespan) this may increase further to 26 cm above the 1981 – 2000 baseline.
- 14.7.6 The direct effect of a relatively small increase in sea level rise within the region of 8 – 11 cm prior to and throughout the construction phase of the Proposed Development is an extension and / or shift in the distribution of intertidal habitats which are used as nursery grounds by several fish species. As there is not predicted to be any significant decline in the availability of functional habitats, effects to fisheries or fish ecology are likely to be limited and not significant.
- 14.7.7 Sea temperature change projections are more variable and less specific to the Teesside region. Under RCP 8.5 a rise in global sea surface temperatures of 1.5°C by 2050 is predicted, increasing to a 3.2°C rise by 2100 relative to 1870 – 1899 temperatures. In UK waters, mean annual sea temperatures have risen by 0.8°C since 1870 and have continued to show consistent warming trends since the 1970s onwards (Genner *et al.*, 2017). According to Lowe *et al.* (2009), the seas around the UK are projected to be 1.5 – 4 °C warmer by 2100.
- 14.7.8 Rapid responses to climate change are likely to be observed for fish species that rely on planktonic food sources owing to climate induced changes to phytoplankton and zooplankton productivity and distribution (Rijnsdorp *et al.*, 2009). Furthermore, species such as mackerel, herring, and sardines (*Sardina* sp.), which are mobile and wide-ranging pelagic species, are expected to have the greatest distributional variations, although changes in the distribution and behaviour of demersal fish have also been reported (Mieszkowska *et al.*, 2009).
- 14.7.9 It is predicted that the distribution of Lusitanian species (those with an affinity for warm water) in UK waters, such as sand smelt and horse mackerel, will shift further north around the coast due to increasing sea temperatures (Heath *et al.*, 2012). Additionally, the optimal habitat range of boreal species (those with an affinity for cold water), such as cod and herring, is also likely to shift northwards or into deeper waters (Heath *et al.*, 2012).

⁵ This baseline has been selected as it provides projections for 20-year time periods (e.g. 2020 – 2039).

- 14.7.10 Species which don't migrate, for instance schooling pelagic species, like the lesser sandeel, are likely to be affected more locally, with potential impacts to abundance (Frederiksen *et al.*, 2006). It is predicted that changes to sea water temperatures may alter the spawning period of certain species, such as whiting, sole, and sprat, who have an energy-income breeding strategy where spawning period is temperature dependent (Heath *et al.*, 2012).
- 14.7.11 In the short-term (i.e. between the time of writing in 2020 and beginning of construction in Q3 2022), climate change is expected to have a negligible impact on fish. However, in the medium-term and long-term (i.e. commencement of operation in 2026 and during the operational lifetime of the Proposed Development, respectively), climate change may have the following impacts:
- Increased frequency and severity of droughts reducing water levels or drying-out of watercourses may impact migratory fish prohibiting access to upstream spawning habitats;
 - Increased frequency and severity of flooding leading to the mortality of fish adults, juveniles, and eggs;
 - Shift in north-south and onshore-offshore distributions of species and functional habitats due to increasing sea temperatures; and
 - Changes in the timing of spawning and other lifecycle characteristics due to increasing sea temperatures.
- 14.7.12 Overall, it is difficult to say with any certainty the potential magnitude of changes to fisheries and fish ecology within the study area prior to and during construction and operation of the Proposed Development. However, it is unlikely that significant changes in baseline conditions would be detectable above natural variability in the short and medium-term but may be observed in the long-term.

14.8 Summary of Findings

- 14.8.1 The Tees river and estuary is an important water body for the transit of a variety of diadromous fish species which make seasonal migrations between the sea and riverine environments. The species known to migrate through the study area include salmon, sea trout, European eel, river lamprey and sea lamprey, all of which have been identified as part of the Tees Valley BAP. The River Tees is particularly important for salmon and sea trout which both regularly use the Tees catchment area. Furthermore, the River Tees is designated as one of 64 'principal salmon rivers' in England and Wales and has its own 'Salmon Action Plan'.
- 14.8.2 A range of estuarine and marine fish and shellfish species have been recorded in the study area, most notably in the lower reaches of the River Tees. The taxa identified represent a mixed demersal and pelagic fish assemblage which is typical of the central North Sea. The highest abundances were of the species sprat, herring and lesser sandeel, which were caught in the intertidal areas of the lower reaches of the River Tees. Cod, whiting and plaice also represented a high proportion of the estuarine



and marine fish caught. Moreover, the study area has been identified as located within the nursery grounds of the following species: herring, sprat, cod, whiting, plaice, Nephrops, lemon sole and spurdog. The Proposed Development is also found within the spawning areas of lemon sole and Nephrops. Spawning of sprat takes place from May to August, peaking in May to June, whilst spawning of herring occurs predominantly in late summer (although spawning can happen at different times). It is therefore expected that the highest abundance of juvenile fish in the intertidal will be in summer and early autumn. Mackerel was not identified as being a key risk species, with only one individual having been recorded in the River Tees in Environment Agency TraC surveys.

- 14.8.3 The key migratory periods for the diadromous fish species utilising the River Tees have been identified, which for salmon migrating as smolt down rivers to oceans, occurs in spring to early summer. The return of salmon as adults arises between June to August and October to December, whilst sea trout have a similar migratory period. The significant migratory periods of European eel take place in spring and continue until early autumn, when eels transition upstream into freshwater, and then returning to spawning grounds between August and December. River lamprey typically migrate into freshwater from October to December and continues to do so through winter into early spring. The river lamprey then migrates back downstream from July to September. Sea lamprey normally migrates into freshwater in April and May as adults, whilst the migration to sea can vary from river to river, although the metamorphosis of larvae into adults following spawning, occurs in July and September.
- 14.8.4 Taking into consideration climate change predictions prior to and during the construction and operational phase of the Proposed Development, it is anticipated that the fish and shellfish baseline is likely to evolve. However, it is unlikely that significant changes in baseline conditions would be detectable above natural variability.



14.9 References

Aerestrup, K., Økland, F., Hansen, M.M., Righton, D., Gargan, P., Castonguay, M., Bernatchez, L., Howey, P., Sparholt, H., Pedersen, M.I. and McKinley, R.S. (2009). Oceanic spawning migration of the European eel (*Anguilla anguilla*). *Science*, 325(5948), 1660 – 1660.

Atema, J. (1986). Review of sexual selection and chemical communication in the lobster *Homarus americanus*. *Canadian Journal of Fisheries and Aquatic Science*. 43, 2283 – 2390.

Bannister, R.C.A., Addison, J.T., and Lovewell, S.R.J. (1994). Growth, movement, recapture rate and survival of hatchery-reared lobsters (*Homarus gammarus* Linnaeus, 1758) released into the wild on the English East Coast. *Crustaceana*. 67, 156 – 172.

Barnes, M. (2008). *Merlangius merlangus*. Whiting. *Marine Life Information Network: Biology and Sensitivity Key Information Sub-programme* [Online]. Available at: <http://www.marlin.ac.uk/speciesinformation.php?speciesID=3794> [Accessed: 16 September 2019].

Behrmann-Godel, J. and Eckmann, R. (2003). A preliminary telemetry study of the migration of silver European eel (*Anguilla anguilla* L.) in the River Mosel, Germany. *Ecology of Freshwater Fish*, 12(3), 196 – 202.

Blaxter, J.H.S., and Hunter, J.R. (1982). The biology of the clupeoid fishes. *Advances in Marine Biology*, 20, 1 – 223.

Braber, L., and Groot, S.J. de. (1973). The food of five flatfish species (Pleuronectiformes) in the southern North Sea. *Netherlands Journal of Sea Research*, 6(1-2), 163 – 172.

Brown and May Marine (2014). Dogger Bank Teesside A and B, Environmental Statement Chapter 13 Appendix A Fish and Shellfish Ecology Technical Report. Application Reference 6.13.1.

Callaway, R., Alsvåg, J., De Boois, I., Cotter, J., Ford, A., Hinz, H., Jennings, S., Kröncke, I., Lancaster, J., Piet, G. and Prince, P. (2002). Diversity and community structure of epibenthic invertebrates and fish in the North Sea. *ICES Journal of Marine Science*. 59(6), 1199 – 1214.

Centre for Environment, Fisheries and Aquaculture Science, Environment Agency, and Natural Resources Wales. (2017a). *Edible crab (Cancer pagurus)*, *Cefas Stock Status Report 2017* [Online]. Available at: https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/722904/Cefas_Crab_Stock_Assessment_2017.pdf [Accessed: 23 September 2019].

Centre for Environment, Fisheries and Aquaculture Science, Environment Agency, and Natural Resources Wales. (2017b). *Lobster (Homarus gammarus)*, *Cefas Stock Status Report 2017* [Online]. Available at: https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/722904/Cefas_Crab_Stock_Assessment_2017.pdf



[/attachment_data/file/722748/2017_Lobster_assessments.pdf](#) [Accessed: 25 September 2019].

Centre for Environment, Fisheries and Aquaculture Science, Environment Agency, and Natural Resources Wales. (2019). *Salmon Stocks and Fisheries in England and Wales in 2018* [Online]. Available at: https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/810720/SalmonReport-2018-assessment_final.pdf [Accessed: 03 December 2019].

Chadwick, S., Knights, B., Thorley, J.L. and Bark, A. (2007). A long-term study of population characteristics and downstream migrations of the European eel *Anguilla anguilla* (L.) and the effects of a migration barrier in the Girnock Burn, north-east Scotland. *Journal of Fish Biology*, 70(5), 1535 – 1553.

Cohen, D.M., Inada, T., Iwamoto, T. and Scialabba, N. (1990). Gadiform fishes of the world. *FAO Fisheries Synopsis*, 10(125).

Cotter, A.J.R., Course, G., Ashworth, J., Forster, R., Enver, R., Goad, D., Bush, R., Shaw, S., Mainprize, R., and Garshelis, D.L. (2004). Summary of commercial marine fishing by English and Welsh vessels over 10 metres LOA for 2004. *Science Series Technical Report* no.134.

Coull, K.A. Johnstone, R. and Rogers, S.I. (1998). *Fisheries Sensitivity Maps in British Waters*. London: UKOOA Ltd.

Cowx, I.G. and Fraser, D. (2003). Monitoring the Atlantic Salmon, Conserving Natura 2000 Rivers Monitoring Series No. 7. *English Nature*.

Daan, N. (1973). A quantitative analysis of the food intake of North Sea cod, *Gadus morhua*. *Netherlands Journal of Sea Research*, 6(4), 479 – 517.

Daan, N., Rijnsdorp, A.D., and Van Overbeeke, G.R. (1985). Predation by North Sea herring *Clupea harengus* on eggs of plaice *Pleuronectes platessa* and cod *Gadus morhua*. *Transactions of the American Fisheries Society*, 114, 499 – 506.

De Clerck, R. and Buseyne, D. (1989). *On the feeding of plaice (Pleuronectes platessa L.) in the southern North Sea*. Copenhagen: International Council for the Exploration of the Sea (ICES).

Degerman, E., Leonardsson, K. and Lundqvist, H. (2012). Coastal migrations, temporary use of neighbouring rivers, and growth of sea trout (*Salmo trutta*) from nine northern Baltic Sea rivers. *ICES Journal of Marine Science*, 69(6), 971 – 980.

Dempsey, C.H. and Bamber, R.N. (1983). Spawning of herring (*Clupea harengus* L.) in the Blackwater Estuary, spring 1979. *ICES Journal of Marine Science*, 41(1), 85 – 92.

Dipper, F. (2001). *British sea fishes*. Unknown: Underwater World.



Eaton, D.R., Brown, J., Addison, J.T., Milligan, S.P. and Fernand, L.J. (2003). Edible crab (*Cancer pagurus*) larvae surveys off the east coast of England: implications for stock structure. *Fisheries Research*, 65(1-3), 191 – 199.

Ellis, J.R. Milligan, S.P. Readdy, L. Taylor, N. and Brown, M.J. (2012). *Spawning and nursery grounds of selected fish species in UK waters*. Science Series Technical Report. Lowestoft: Cefas. 147, 56 pp.

Entec UK Limited. (2011). *Teesside Offshore Wind Farm FEPA Monitoring*. Annual Fish Survey Report. 55 pp.

Environment Agency (2009). *River Tees Salmon Action Plan Review*. APEM Scientific Report EA 410441.

Environment Agency (2013). *River Tees fish counts* [Online]. Available at: <https://webarchive.nationalarchives.gov.uk/20140328235559/http://www.environment-agency.gov.uk/research/library/publications/138616.aspx> [Accessed: 13 September 2019].

Environment Agency (2017). *Salmonid and Freshwater Fisheries Statistics for England and Wales (2017). Including declared catches for salmon, sea trout, eels, smelt and lamprey by rods, nets and other instruments*. Version 2, 25th April 2019.

Environment Agency (2018). *The current state of salmon stocks* [Online]. 57 pp. Available at: https://consult.environment-agency.gov.uk/fisheries/proposed-national-salmon-byelaws/results/appendix5_thecurrentstateofsalmonstocks_mar2018.pdf

Environment Agency (2019a). *TraC Fish Counts for all Species for all Estuaries and all years* [Online]. Available at: <https://data.gov.uk/dataset/41308817-191b-459d-aa39-788f74c76623/trac-fish-counts-for-all-species-for-all-estuaries-and-all-years> [Accessed: 17 September 2019].

Environment Agency (2019b). *Fish Counting at Tees Barrage* [Online]. Available at: https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/825747/Tees_monthly_counts.pdf [Accessed: 18 September 2019].

Environment Agency (2019c). *Freshwater Fish Counts for all Species, all Areas and all Years* [Online]. Available at: <https://data.gov.uk/dataset/f49b8e4b-8673-498e-bead-98e6847831c6/freshwater-fish-counts-for-all-species-all-areas-and-all-years> [Accessed: 17 January 2020].

Farmer, A.S.D. (1974). Reproduction in *Nephrops norvegicus* (Decapoda: nephropidae). *Journal of Zoology*, 174(2), 61 – 183.

Frederiksen, M., Edwards, M., Richardson, A.J., Halliday, N.C. and Wanless, S. (2006). From plankton to top predators: bottom-up control of a marine food web across four trophic levels. *Journal of Animal Ecology*, 75(6), 1259 – 1268.



Gargan, P.G., Roche, W.K., Forde, G.P. and Ferguson, A. (2006). Characteristics of the sea trout (*Salmo trutta* L.) stocks from the Owengowla and Invermore fisheries, Connemara, Western Ireland, and recent trends in marine survival. *Sea trout: biology, conservation and management*. Blackwell, Oxford, 60 – 75.

Gauld, N.R., Campbell, R.N.B. and Lucas, M.C. (2013). Reduced flow impacts salmonid smolt emigration in a river with low-head weirs. *Science of the total environment*, 458, 435 – 443.

Green, E. (2017). Tern diet in the UK and Ireland: a review of key prey species and potential impacts of climate change. RSPB.

Greenstreet, S.P.R., Holland, G.J., Guirey, E.J., Armstrong, A., Fraser, H.M., and Gibb, I.M. (2010). Combining hydroacoustic seabed survey and grab sampling techniques to assess “local” sandeel population abundance. *ICES Journal of Marine Science*. 67, 971 – 984.

Harris, G. (2017). Sea Trout – Science and Management. *Proceedings of the 2nd International Sea Trout Symposium held in Dundalk, Republic of Ireland, on 20 – 22 October 2015*. Inland Fisheries Ireland.

Heath, M.R., Neat, F.C., Pinnegar, J.K., Reid, D.G., Sims, D.W. and Wright, P.J. (2012). Review of climate change impacts on marine fish and shellfish around the UK and Ireland. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 22(3), 337 – 367.

Heessen, H.J. and Rijnsdorp, A.D. (1989). Investigations on egg production and mortality of cod (*Gadus morhua* L.) and plaice (*Pleuronectes platessa* L.) in the southern and eastern North Sea in 1987 and 1988. *Conseil Permanent International pour l'Exploration de la Mer*, 191, 15 – 20.

Heessen, H.J. (1993). The distribution of cod (*Gadus morhua*) in the North Sea. *NAFO Scientific Council Studies*, 18, 7 pp.

Heesscn, H.J.L. and Daan, N. (1994). Cod distribution and temperature in the North Sea. *ICES Marine Science Symposia*. 198, 244 – 253.

Hinrichsen, H.H. Kraus, G. Voss, R., Stepputtis, D. and Baumann, H. (2005). The general distribution pattern and mixing probability of Baltic sprat juvenile populations. *Journal of Marine Systems*. 58, 52 – 66.

Hislop, J.R.G. and Hall, W.B. (1974). The fecundity of whiting, *Merlangius merlangus* (L.) in the North Sea, the Minch and at Iceland. *ICES Journal of Marine Science*, 36(1), 42 – 49.

Hislop, J.R.G., Robb, A.P., Bell, M.A. and Armstrong, D.W. (1991). The diet and food consumption of whiting (*Merlangius merlangus*) in the North Sea. *ICES Journal of Marine Science*, 48(2), 139 – 156.

Holland, G.J., Greenstreet, S.P.R., Gibb, I.M., Fraser H.M., and Robertson, M.R. (2005). Identifying Sandeel *Ammodytes marinus* sediment habitat preferences in the marine environment. *Marine Ecology Progress Series*. 303, 269 – 282.



Howard, A.E. (1982). The distribution and behaviour of ovigerous edible crabs (*Cancer pagurus*), and consequent sampling bias. *Journal du Conseil International pour l'Exploration de la Mer*, 40, 259 – 261.

Howard, F.G. (1989). *The Norway Lobster*. Scottish Fisheries Information Pamphlet Number 7 1989 (Second Edition) ISSN 03099105.

International Council for the Exploration of the Sea (ICES) (1965). *Report of the working group on sole*. ICES Cooperative Research Report 5: 1 - 126

International Council for the Exploration of the Sea (ICES) (1999). *Report of the working group on the assessment of Mackerel, Horse Mackerel, Sardine and Anchovy (WGMHSA)*. ICES CM2000/ACFM:5.

International Council for the Exploration of the Sea (ICES) (2005). *Report of the Study Group of Multispecies Assessment in the North Sea (SGMSNS)*, 5 -8 April 2005, ICES Headquarters. ICES CM 2005/D:06, pp. 163

International Council for the Exploration of the Sea (ICES) (2006a). *ICES FishMap, Species Factsheet – herring* [Online]. Available at: <http://www.ices.dk/explore-us/projects/EU-RFP/EU%20Repository/ICES%20FishMap/ICES%20FishMap%20species%20factsheet-herring.pdf> [Accessed: 7 October 2019].

International Council for the Exploration of the Sea (ICES) (2006b). *ICES FishMap, Species Factsheet – sprat* [Online]. Available at: <http://www.ices.dk/explore-us/projects/EU-RFP/EU%20Repository/ICES%20FishMap/ICES%20FishMap%20species%20factsheet-sprat.pdf> [Accessed: 7 October 2019].

International Council for the Exploration of the Sea (ICES) (2006c). *ICES FishMap, Species Factsheet – mackerel* [Online]. Available at: <http://www.ices.dk/explore-us/projects/EU-RFP/EU%20Repository/ICES%20FishMap/ICES%20FishMap%20species%20factsheet-mackerel.pdf> [Accessed: 7 October 2019].

International Council for the Exploration of the Sea (ICES) (2006d). *ICES FishMap, Species Factsheet – cod* [Online]. Available at: <http://www.ices.dk/explore-us/projects/EU-RFP/EU%20Repository/ICES%20FishMap/ICES%20FishMap%20species%20factsheet-cod.pdf> [Accessed: 7 October 2019].

International Council for the Exploration of the Sea (ICES) (2006e). *ICES FishMap, Species Factsheet – sole* [Online]. Available at: <http://www.ices.dk/explore-us/projects/EU-RFP/EU%20Repository/ICES%20FishMap/ICES%20FishMap%20species%20factsheet-sole.pdf> [Accessed: 7 October 2019].

International Council for the Exploration of the Sea (ICES) (2006f). *ICES FishMap, Species Factsheet – plaice* [Online]. Available at: <http://www.ices.dk/explore-us/projects/EU-RFP/EU%20Repository/ICES%20FishMap/ICES%20FishMap%20species%20factsheet-plaice.pdf> [Accessed: 7 October 2019].



International Council for the Exploration of the Sea (ICES) (2010). *Report of the ICES/EIFAC Working Group on Eels (WGEEL)*. ICES Document CM 2010/ACFM: 18

International Council for the Exploration of the Sea (ICES) (2008). *Report of the working group on multispecies assessment methods (WGSAM)*. ICES CM 2008/RMC:06.

International Council for the Exploration of the Sea (ICES) (2011). *Report of the Working Group on Widely Distributed Stocks (WGWIDE)*. ICES CM 2011/ACOM:15

International Council for the Exploration of the Sea (ICES) (2016). *Report of the Working Group on the Biology and Life History of Crabs (WGCRAB)*. [Online]. Available at:

<http://www.ices.dk/sites/pub/Publication%20Reports/Expert%20Group%20Report/SSGEPD/2016/01%20WGCRAB%20-%20Report%20of%20the%20Working%20Group%20on%20the%20Biology%20and%20Life%20History%20of%20Crabs.pdf> [Accessed: 23 September 2019].

International Council for the Exploration of the Sea (ICES) (2019a). *Sprat (Sprattus sprattus) in Division 3.a and Subarea 4 (Skagerrak, Kattegat, and North Sea)* [Online]. Available at:

<https://www.ices.dk/sites/pub/Publication%20Reports/Advice/2019/2019/spr.27.3a4.pdf> [Accessed: 24 September 2019].

International Council for the Exploration of the Sea (ICES) (2019b). *Norway special request for revised 2019 advice on mackerel (Scomber scombrus) in subareas 1–8 and 14, and in Division 9.a (the Northeast Atlantic and adjacent waters)* [Online]. Available at:

http://www.ices.dk/sites/pub/Publication%20Reports/Advice/2019/Special_Request/no.2019.09.pdf [Accessed: 22 September 2019].

International Council for the Exploration of the Sea (ICES) (2019c). *Cod (Gadus morhua) in Subarea 4, Division 7.d, and Subdivision 20 (North Sea, eastern English Channel, Skagerrak)* [Online]. Available at:

<http://ices.dk/sites/pub/Publication%20Reports/Advice/2019/2019/cod.27.47d20.pdf> [Accessed: 21 September 2019].

International Council for the Exploration of the Sea (ICES) (2019d). *Plaice (Pleuronectes platessa) in Subarea 4 (North Sea) and Subdivision 20 (Skagerrak)* [Online]. Available at:

<http://www.ices.dk/sites/pub/Publication%20Reports/Advice/2019/2019/ple.27.420.pdf> [Accessed: 23 September 2019].

International Council for the Exploration of the Sea (ICES) (2019e). *Sandeel (Ammodytes spp.) in divisions 4.a–b, Sandeel Area 4 (northern and central North Sea)* [Online]. Available at:

<http://www.ices.dk/sites/pub/Publication%20Reports/Advice/2019/2019/san.s4.a.4.pdf> [Accessed: 23 September 2019].



International Council for the Exploration of the Sea (ICES) (2019f). *Norway lobster (Nephrops norvegicus) in Division 4.b, Functional Unit 6 (central North Sea, Farn Deep)* [Online]. Available at: <http://www.ices.dk/sites/pub/Publication%20Reports/Advice/2019/2019/nep.f.u.6.pdf> [Accessed: 24 September 2019].

Jansen, T. and Gislason, H. (2011). Temperature affects the timing of spawning and migration of North Sea mackerel. *Continental Shelf Research*, 31(1), 64 – 72.

Joint Nature Conservation Committee (JNCC) (2013a). *European Community Directive on the Conservation of Natural Habitats and of Wild Fauna and Flora (92/43/EEC), Third Report by the United Kingdom under Article 17, on the implementation of the Directive from January 2007 to December 2012 Conservation status assessment for S1095 - Sea lamprey (Petromyzon marinus)* [Online]. Available at: http://archive.jncc.gov.uk/pdf/Article17Consult_20131010/S1095_UK.pdf [Accessed: 22 September 2011].

Joint Nature Conservation Committee (JNCC) (2013b). *European Community Directive on the Conservation of Natural Habitats and of Wild Fauna and Flora (92/43/EEC), Third Report by the United Kingdom under Article 17, on the implementation of the Directive from January 2007 to December 2012 Conservation status assessment for S1099 - River lamprey (Lampetra fluviatilis)* [Online]. Available at: http://archive.jncc.gov.uk/pdf/Article17Consult_20131010/S1095_UK.pdf [Accessed: 22 September 2011].

Kelly, F. L. and King, J. J. (2001). A review of the ecology and distribution of three lamprey species, *Lampetra fluviatilis* (L.), *Lampetra planeri* (L.), and *Petromyzon marinus* (L.): a context for conservation and biodiversity considerations in Ireland. *Biology and Environment: Proceedings of the Royal Irish Academy*. 101(3), 165 – 185.

King, R.A., Hillman, R., Elsmere, P., Stockley, B. and Stevens, J.R. (2016). Investigating patterns of straying and mixed stock exploitation of sea trout, *Salmo trutta*, in rivers sharing an estuary in south-west England. *Fisheries Management and Ecology*, 23(5), 376 – 389.

Kjesbu, O.S., Kryvi, H., Sundby, S. and Solemdal, P. (1992). Buoyancy variations in eggs of Atlantic cod (*Gadus morhua* L.) in relation to chorion thickness and egg size: theory and observations. *Journal of Fish Biology*, 41(4), 581 – 599.

Kruuk, H. (1963). Diurnal periodicity in the activity of the common sole, *Solea vulgaris* Quensel. *Netherlands Journal of Sea Research* 2(1), 1 – 28.

Kuipers, B.R. (1977). On the Ecology of Juvenile Plaice on A Tidal Flat in the Wadden Sea. *Netherlands Journal of Sea Research*, 11, 56 – 91.

Lancaster, J., Taylor, E., Lowe, A., McCallum, S. (2011). *Teesside Windfarm Ltd, Teesside Offshore Wind Farm FEPA Monitoring, Annual Fish Survey Report*. Entek UK Ltd.



Last, J.M. (1978). The food of three species of gadoid larvae in the eastern English Channel and southern North Sea. *Marine Biology*, 48(4), 377 – 386.

Last, J.M. (1989). The food of herring, *Clupea harengus*, in the North Sea, 1983-1986. *Journal of Fish Biology*, 34, 489 – 501.

Laughton, R. and Burns, S. (2003). *Assessment of sea lamprey distribution and abundance in the River Spey: Phase III*. Scottish Natural Heritage Commissioned Report No. 043 (ROAME No. F02AC604).

Limpenny, S.E., Barrio Froján, C., Cotterill, C., Foster-Smith, R.L., Pearce, B., Tizzard, L., Limpenny, D.L., Long, D., Walmsley, S., Kirby, S., Baker, K., Meadows, W.J., Rees, J., Hill, J., Wilson, C., Leivers, M., Churchley, S.,

Macer, C.T. (1966). Sand eels (*Ammodytidae*) in the south western North Sea: their biology and fishery. *Fish Investment London Series II*. 24, 1 – 55.

Macer, C.T. (1976). *Observations on the maturity and fecundity of mackerel (Scomber scombrus, L.)*. ICES CM.

Maes, J. and Ollevier, F. (2002). Size structure and feeding dynamics in estuarine clupeid fish schools: field evidence for the school trap hypothesis. *Aquatic Living Resources*. 15, 211 – 216.

Maitland, P. S. (2003). *Ecology of the River Brook and Sea Lamprey. Conserving Natura 2000 Rivers Ecology Series No. 5*. Peterborough: English Nature.

Malcolm. I.A., Godfrey, J. and Youngson. A.F. (2010). Review of migratory routes and behaviour of Atlantic salmon, sea trout and European eel in Scotland's coastal environment: implications for the development of marine renewables. *Scottish Marine and Freshwater Science* Vol 1 No 14.

Marine Management Organisation (MMO) (2016). *North East Marine Plan* [Online]. Available at: <https://www.gov.uk/government/collections/north-east-marine-plan#north-east-marine-plan-areas> [Accessed: 9 September 2019].

Marine Management Organisation (MMO) (2018). *UK sea fisheries annual statistics report 2017* [Online]. Available at: <https://www.gov.uk/government/statistics/uk-sea-fisheries-annual-statistics-report-2017> [Accessed: 6 September 2019].

MarineSpace Ltd, ABPmer Ltd, ERM Ltd, Fugro EMU Ltd and Marine Ecological Surveys Ltd (2013). *Environmental Effect Pathways between Marine Aggregate Application Areas and Sandeel Habitat: Regional Cumulative Impact Assessments*. Version 1.0. A report for the British Marine Aggregates Producers Association.

McCormick, S.D., Hansen, L.P., Quinn, T.P. and Saunders, R.L. (1998). Movement, migration, and smolting of Atlantic salmon (*Salmo salar*). *Canadian Journal of Fisheries and Aquatic Sciences*, 55(S1), 77 – 92.

McEvoy, L.A. and McEvoy, J. (1992). Multiple spawning in several commercial fish species and its consequences for fisheries management, cultivation and experimentation. *Journal of Fish Biology*, 41, 125 – 136.



Mieszowska, N., Genner, M.J., Hawkins, S.J. and Sims, D.W. (2009). Effects of climate change and commercial fishing on Atlantic cod *Gadus morhua*. *Advances in marine biology*, 56, 213 – 273.

Milligan, S. P. (1986). *Recent studies on the spawning of sprat (Sprattus sprattus L.) in the English Channel*. Fisheries Research Technical Report no. 83.

Moore, J. A. Hartel, K. E. Craddock, J. E. and Galbraith, J. K. (2003). An annotated list of deepwater fishes from off the New England region, with new area records. *Northeastern Naturalist*, 10(2), 159 – 248.

Moore, A., and Potter, E.C.E. (2014). *Provisional Assessment of the River Tees Barrage Fish Passage*. CEFAS.

Murua, H. and Saborido-Rey, F. (2003). Female reproductive strategies of marine fish species of the North Atlantic. *Journal of Northwest Atlantic Fisheries Science*, 33, 23 – 31.

Natural England (2010). *The Dee Estuary European Marine Site. Natural England & the Countryside Council for Wales advice given under Regulation 33(2) of the Conservation (Natural Habitats &c.) Regulations 1994*.

Neal, K.J. and Wilson, E. (2008). *Cancer pagurus* Edible crab. In Tyler-Walters H. and Hiscock K. (eds) *Marine Life Information Network: Biology and Sensitivity Key Information Reviews*. Plymouth: Marine Biological Association of the United Kingdom [Online]. Available at: <https://www.marlin.ac.uk/species/detail/1179> [Accessed: 26 September 2019].

Nissling, A., Muller, A. and Hinrichsen, H.H. (2003). Specific gravity and vertical distribution of sprat (*Sprattus sprattus*) eggs in the Baltic Sea. *Fisheries Biology* 63, 280 – 299.

North Atlantic Salmon Conservation Organisation (NASCO) (2012). *The Atlantic Salmon* [Online]. Available at: <http://www.nasco.int/atlanticsalmon.html> [Accessed: 26 September 2019].

Pawson, M.G. (1995). *Biogeographical identification of English Channel fish and shellfish stocks*. Fisheries Research Technical Report. Lowestoft: Directorate of Fisheries Research, 99, 1 – 72.

Potter, E.C.E. and Dare, P.J. (2003). Research on migratory salmonids, eel and freshwater fish stocks and fisheries. *Science series technical report-centre for environment fisheries and aquaculture science*.

Precision Marine Survey Ltd (2014). *Dogger Bank Teesside A and B, Environmental Statement Chapter 13 Appendix C Nearshore Fish and Shellfish Surveys*. Application Reference 6.13.3.

Reid, J.E. and Chaput, G. (2012). Spawning history influence on fecundity, egg size, and egg survival of Atlantic salmon (*Salmo salar*) from the Miramichi River, New Brunswick, Canada. *ICES Journal of Marine Science*, 69(9), 1678 – 1685.



Reiss, H., Degraer, S., Duineveld, G.C., Kröncke, I., Aldridge, J., Craeymeersch, J.A., Eggleton, J.D., Hillewaert, H., Lavaleye, M.S., Moll, A. and Pohlmann, T. (2009). Spatial patterns of infauna, epifauna, and demersal fish communities in the North Sea. *ICES Journal of Marine Science*. 67(2), 278 – 293.

Righton, D., Westerberg, H., Feunteun, E., Økland, F., Gargan, P., Amilhat, E., Metcalfe, J., Lobon-Cervia, J., Sjöberg, N., Simon, J. and Acou, A. (2016). Empirical observations of the spawning migration of European eels: The long and dangerous road to the Sargasso Sea. *Science Advances*, 2(10), 15 pp.

Rijnsdorp, A.D., Van Beek, F.A., Flatman, S., Millner, R.M., Riley, J.D., Giret, M. and De Clerck, R. (1992). Recruitment of sole stocks, *Solea solea* (L.), in the Northeast Atlantic. *Netherlands Journal of Sea Research*, 29(1-3), 173 – 192.

Rijnsdorp, A.D., Peck, M.A., Engelhard, G.H., Möllmann, C. and Pinnegar, J.K. (2009). Resolving the effect of climate change on fish populations. *ICES Journal of Marine Science*, 66(7), 1570 – 1583.

Russel, F.S. (1976). The Planktonic Stages of British Marine Fishes, Academic Pres Inc. Ltd., London, 524 pp.

Russell, J., Birchenough, A.C., Green, S.L., and Law, R.J. (2011). *The East Coast Regional Environmental Characterisation*. Cefas Open report 08/04. 287 pp.

Ryland, J.S., 1964. The feeding of plaice and sand-eel larvae in the southern North Sea. *Journal of the Marine Biological Association of the United Kingdom*, 44(2), 343 – 364.

Special Committee on Seals (SCOS) (2018). *Scientific Advice on Matters Related to the Management of Seal Populations: 2018*. Sea Mammal Research Unit [Online]. Available at: <http://www.smru.st-andrews.ac.uk/research-policy/scos/> [Accessed: 6 September 2019].

Sheridan, S., Turnpenny, A., Horsfield, D., Bamford, D., Bayliss, B., Coates, S., Dolben, I., Frear, P., Hazard, E., Tavner, I. and Trudgill, N. (2013). Screening at intakes and outfalls: measures to protect eel (*Anguilla anguilla*). *International Fish Screening Techniques*, 71. 17 – 30.

Shrimpton, J.M. (2012). Seawater to freshwater transitions in diadromous fishes. In *Fish physiology*. Academic Press. 32, 327 – 393.

Tappin, D R, Pearce, B, Fitch, S, Dove, D, Gearey, B, Hill, J M, Chambers, C, Bates, R, Pinnion, J, Diaz Doce, D, Green, M, Gallyot, J, Georgiou, L, Brutto, D, Marzialetti, S, Hopla, E, Ramsay, E, and Fielding, H. (2011). *The Humber Regional Environmental Characterisation*. British Geological Survey Open Report OR/10/54. 357 pp.

Teal, L.R. (2011). *The North Sea fish community: past, present and future*. Background document for the 2011 National Nature Outlook.



ter Hofstede, R. H., Winter, H. V. and Bos, O. G. (2008). *Distribution of fish species for the generic Appropriate Assessment for the construction of offshore wind farms*. IMARES Report C050/08.

Tesch, F.-W. (2003). *The eel*. 5th edition. Oxford: Blackwell Science, pp. 416

Thorstad, E.B., Whoriskey, F., Uglem, I., Moore, A., Rikardsen, A.H. and Finstad, B. (2012). A critical life stage of the Atlantic salmon *Salmo salar*: behaviour and survival during the smolt and initial post-smolt migration. *Journal of Fish Biology*, 81(2), 500 – 542.

Trippel, E.A. (1998). Egg size and viability and seasonal offspring production of young Atlantic cod. *Transactions of the American Fisheries Society*, 127(3), 339 – 359.

UK Biodiversity Action Plan. (2012). *Priority habitats and species in the Tees Valley – Update January 2012* [Online]. Available at: <https://teesvalleynaturepartnership.org.uk/wp-content/uploads/2012/11/Tees-Valley-priority-habitats-and-species-updated-5-jan-2012-pdf> [Accessed: 12 September 2019].

van der Kooij, J., Scott, B.E., and Mackinson, S. (2008). The effects of environmental factors on daytime sandeel distribution and abundance on the Dogger Bank. *Journal of Sea Research*. 60(3), 201 – 209.

Waldman, J., Grunwald, C. and Wirgin, I. (2008). Sea lamprey *Petromyzon marinus*: an exception to the rule of homing in anadromous fishes. *Biology letters*, 4(6), 659 – 662.

Wheeler, A. (1978). *Key to the fishes of Northern Europe*. London: Frederik Warne (Publishers) Ltd.

Whitehead, P.J.P. (1986). *Clupeidae*. In *Fishes of the North-eastern Atlantic and the Mediterranean Volume I* (Whitehead, P.J.P., Bauchot, M.-L., Hureau, J.-C., Nielsen, J., and Tortonese, E., eds.) UNESCO, Paris, 268 – 281.

Wilson, E. (2008). *Homarus gammarus Common lobster*. In: Tyler-Walters H. and Hiscock K. (eds) *Marine Life Information Network: Biology and Sensitivity Key Information Reviews*, Plymouth: Marine Biological Association of the United Kingdom [Online]. Available at: <https://www.marlin.ac.uk/species/detail/1171> [Accessed: 26 September 2019].

Wilson, L.J. and Hammond, P.S. (2016). *Harbour seal diet composition and diversity*. Marine Mammal Scientific Support Research Programme MMSS/001/11 CSD 3.2. Report to the Scottish Government.

Winslade P. (1971). *Behavioural and embryological studies on the lesser sand eel Ammodytes marinus (Raftt)*. PhD Thesis. University East Anglia.

Zheng, X., Pierce, G.J., Reid, D.G. and Jolliffe, I.T., 2002. Does the North Atlantic current affect spatial distribution of whiting? Testing environmental hypotheses using statistical and GIS techniques. *ICES Journal of Marine Science*, 59(2), 239 – 253.

Annex A – Temporal variations in the type of gear used during the Environmental TraC surveys

Lower Reaches

Table A1- 1: Temporal variations in the type of gear used during the Environmental TraC surveys in the lower reaches of the River Tees. From 1985 to 2004 beam trawling (2.4 m) was the only sampling method used.

Sampling Year	Sampling Season (Month)	Seine Netting	Beam Trawling – 1.5 m	Beam Trawling – 2.4 m	Otter Trawling
2004	Autumn (October)	X		X	
2005	Summer (June)	X			
	Autumn (September, October)	X		X	
2006	Summer (July)	X			
	Autumn (October)	X		X	
2007	Spring (May)	X			
	Autumn (September, November)	X		X	
	Winter (December)			X	
2008	Summer (June)	X			
	Autumn (October)	X		X	
2009	Summer (June)	X			
	Autumn (October)	X		X	
2010	Summer (June)	X			
	Autumn (September)	X			
2011	Spring (May)	X	X		
	Autumn (September)	X	X		
2012	Spring (May)	X	X		
	Autumn (September)	X	X		
2013	Spring (May)	X			

Sampling Year	Sampling Season (Month)	Seine Netting	Beam Trawling – 1.5 m	Beam Trawling – 2.4 m	Otter Trawling
	Autumn (September)	X	X		
2014	Summer (June)	X	X		
	Autumn (October)				X
2015	Summer (June)	X	X		
	Autumn (September)	X	X		X
2016	Autumn (October, November)	X	X		X
2017	Spring (May)	X	X		
	Autumn (September, November)	X	X		X
2018	Autumn (October)	X	X		X

Middle Reaches

Table A1- 2: Temporal variations in the type of gear used during the Environmental TraC surveys in the middle reaches of the River Tees. From 1981 to 2002 beam trawling (2.4 m) was the only sampling method used.

Sampling Year	Sampling Season (Month)	Seine Netting	Beam Trawling – 1.5 m	Beam Trawling – 2.4 m	Otter Trawling
2002	Autumn (October)			X	X
2003	Autumn (October)			X	
2004	Autumn (October)	X		X	
2005	Autumn (September)	X			
2006	Autumn (October)	X			
2007	Spring (May)	X			
	Autumn (September, November)	X		X	
	Winter (December)			X	
2008	Summer (June)	X			
	Autumn (October)	X			
2009	Summer (June)	X			
	Autumn (October)	X			

Sampling Year	Sampling Season (Month)	Seine Netting	Beam Trawling – 1.5 m	Beam Trawling – 2.4 m	Otter Trawling
2010	Summer (June)	X			
	Autumn (September)	X			
2011	Spring (May)	X			
	Autumn (September)	X			
2012	Spring (May)	X			
	Autumn (September)	X			
2013	Spring (May)	X			
	Autumn (September)	X			
2014	Summer (June)	X			
2015	Summer (June)	X			
	Autumn (September)	X			

Upper Reaches

Table A1- 3: Temporal variations in the type of gear used during the Environmental TraC surveys in the upper reaches of the River Tees.

Sampling Year	Sampling Season (Month)	Seine Netting	Beam Trawling – 1.5 m	Beam Trawling – 2.4 m	Otter Trawling
1982	Summer (June)			X	
	Autumn (September)			X	
1983	Spring (March)			X	
	Summer (June)			X	
	Autumn (September)			X	
1984	Spring (April, May)			X	
	Summer (June, July, August)			X	
	Autumn (September)			X	
1985	Spring (April, May)			X	
	Summer (June, July, August)			X	
	Autumn (September)			X	

Sampling Year	Sampling Season (Month)	Seine Netting	Beam Trawling – 1.5 m	Beam Trawling – 2.4 m	Otter Trawling
1986	Autumn (September, November)			X	
1987	Spring (April)			X	
	Summer (June, July)			X	
	Autumn (September, October)			X	
1988	Spring (April)			X	
	Summer (June, August)			X	
	Autumn (October, November)			X	
1989	Spring (May)			X	
	Summer (June)			X	
	Autumn (September)			X	
1990	Winter (February)			X	
	Spring (April, May)			X	
	Summer (July, August)			X	
	Autumn (November)			X	
1991	Winter (January)			X	
	Spring (April, May)			X	
	Summer (July)			X	
	Autumn (September, November)			X	
1992	Winter (February)			X	
	Spring (April)			X	
	Autumn (September)			X	
1996	Winter (February)			X	

Annex B – Full species list and proportions as presented in TraC survey fish species proportion figures.

Table B1- 1: Proportion of fish species from Environment Agency (2019a) TraC surveys in the lower reaches of the River Tees (1985 – 2018). Fish species in the legend have been presented in order of percentage contribution.

Fish species	Latin name	Proportion
Sprat	<i>Sprattus sprattus</i>	41%
Herring	<i>Clupea harengus</i>	15%
Lesser sandeel	<i>Ammodytes tobianus</i>	10%
Plaice	<i>Pleuronectes platessa</i>	9%
Dab	<i>Limanda limanda</i>	7%
Whiting	<i>Merlangius merlangus</i>	5%
Cod	<i>Gadus morhua</i>	3%
Saithe	<i>Pollachius virens</i>	3%
Flounder	<i>Platichthys flesus</i>	2%
Pogge	<i>Agonus cataphractus</i>	2%
Sandeel spp.	<i>Ammodytidae</i>	2%
Sand goby	<i>Pomatoschistus minutus</i>	1%
Corbin's sandeel	<i>Hyperoplus immaculatus</i>	1%
Lesser weever	<i>Echiichthys vipera</i>	<1%
Short-spined sea scorpion	<i>Myoxocephalus scorpius</i>	<1%
Long-spined sea scorpion	<i>Taurulus bubalis</i>	<1%
Greater sandeel	<i>Hyperoplus lanceolatus</i>	<1%
Five-bearded rockling	<i>Ciliata mustela</i>	<1%
Dragonet	<i>Callionymus lyra</i>	<1%
Dover sole	<i>Solea solea</i>	<1%
Snake pipefish	<i>Entelurus aequoreus</i>	<1%
Viviparous blenny	<i>Zoarces viviparus</i>	<1%
Nilson's pipefish	<i>Syngnathus rostellatus</i>	<1%
Three-spined stickleback	<i>Gasterosteus aculeatus</i>	<1%


Fish species	Latin name	Proportion
Grey gurnard	<i>Eutrigla gurnardus</i>	<1%
Butterfish	<i>Pholis gunnellus</i>	<1%
Greater pipefish	<i>Syngnathus acus</i>	<1%
Red gurnard	<i>Aspitrigla cuculus</i>	<1%
European eel	<i>Anguilla anguilla</i>	<1%
Sea trout	<i>Salmo trutta</i>	<1%
Gurnard sp.	<i>Triglidae</i>	<1%
Lemon sole	<i>Microstomus kitt</i>	<1%
Bib	<i>Trisopterus luscus</i>	<1%
Sea bass	<i>Dicentrarchus labrax</i>	<1%
Sea-snail	<i>Liparis liparis</i>	<1%
Turbot	<i>Psetta maxima</i>	<1%
Angler fish	<i>Lophius piscatorius</i>	<1%
Long rough dab	<i>Hippoglossoides platessoides</i>	<1%
Mackerel	<i>Scomber scombrus</i>	<1%
Painted goby	<i>Pomatoschistus pictus</i>	<1%
Poor cod	<i>Trisopterus minutus</i>	<1%
Sand smelt	<i>Atherina presbyter</i>	<1%
Thin lipped grey mullet	<i>Liza ramada</i>	<1%
Thornback ray	<i>Raja clavata</i>	<1%

Table B1- 2: Proportion of fish species from Environment Agency (2019a) TraC surveys in the middle reaches of the River Tees (1981 – 2018). Fish species in the legend have been presented in order of percentage contribution.


Fish species	Latin name	Proportion
Sprat	<i>Sprattus sprattus</i>	31%
Plaice	<i>Pleuronectes platessa</i>	21%
Whiting	<i>Merlangius merlangus</i>	13%
Herring	<i>Clupea harengus</i>	13%
Flounder	<i>Platichthys flesus</i>	9%
Three-spined stickleback	<i>Gasterosteus aculeatus</i>	4%
Dab	<i>Limanda limanda</i>	2%

Fish species	Latin name	Proportion
Sand goby	<i>Pomatoschistus minutus</i>	2%
Cod	<i>Gadus morhua</i>	2%
Common goby	<i>Pomatoschistus microps</i>	1%
Viviparous blenny	<i>Zoarces viviparus</i>	1%
Long-spined sea scorpion	<i>Taurulus bubalis</i>	<1%
Saithe	<i>Pollachius virens</i>	<1%
Pogge	<i>Agonus cataphractus</i>	<1%
Short-spined sea scorpion	<i>Myoxocephalus scorpius</i>	<1%
European eel	<i>Anguilla anguilla</i>	<1%
Red gurnard	<i>Aspitrigla cuculus</i>	<1%
Butterfish	<i>Pholis gunnellus</i>	<1%
Lesser sandeel	<i>Ammodytes tobianus</i>	<1%
5-bearded rockling	<i>Ciliata mustela</i>	<1%
Dragonet	<i>Callionymus lyra</i>	<1%
Sea trout	<i>Salmo trutta</i>	<1%
Bib	<i>Trisopterus luscus</i>	<1%
Snake pipefish	<i>Entelurus aequoreus</i>	<1%
Dover sole	<i>Solea solea</i>	<1%
River lamprey	<i>Lampetra fluviatilis</i>	<1%
Grey gurnard	<i>Eutrigla gurnardus</i>	<1%
Lesser weever	<i>Echiichthys vipera</i>	<1%
Pollack	<i>Pollachius pollachius</i>	<1%
Roach	<i>Rutilus rutilus</i>	<1%
Atlantic salmon	<i>Salmo salar</i>	<1%
Common bream	<i>Abramis brama</i>	<1%
Dace	<i>Leuciscus leuciscus</i>	<1%
Greater pipefish	<i>Syngnathus acus</i>	<1%
Haddock	<i>Melanogrammus aeglefinus</i>	<1%
Sea-snail	<i>Liparis liparis</i>	<1%

Annex C – Freshwater Fish Count NFPD Environment Agency Survey Locations



This drawing has been produced for the use of AECOM's client. It may not be used, modified or relied upon by third parties, except as agreed by AECOM or as required by law. AECOM accepts no responsibility, and denies any liability whatsoever, to any party that uses or relies upon this drawing without AECOM's express written consent. All dimensions are indicative and in metres unless otherwise noted. Do not scale this document.



PROJECT

NET ZERO TEESSIDE

CLIENT

NZT POWER AND NZNS STORAGE

KEY

NFPD Fish Count Data - Atlantic Salmon

Total Count

1 - 50

51 - 100

101 - 150

151 - 200

201 - 250

251 - 350

Tees Mouth

Tees Barrage

TITLE

FIGURE 14B-C1
NFPD FRESHWATER FISH COUNTS
ATLANTIC SALMON

REFERENCE


NZT_200526_FEB_14B-
C1_v2

SHEET NUMBER

1 of 1

DATE

10/06/2020



This drawing has been produced for the use of AECOM's client. It may not be used, modified or relied upon by third parties, except as agreed by AECOM or as required by law. AECOM accepts no responsibility, and denies any liability whatsoever, to any party that uses or relies upon this drawing without AECOM's express written consent. All dimensions are indicative and in metres unless otherwise noted. Do not scale this document.



PROJECT

NET ZERO TEESSIDE

CLIENT

NTZ POWER AND NZNS STORAGE

KEY

NFPD Fish Count Data - Brown/Sea Trout

Total Count

1 - 50

51 - 100

101 - 150

151 - 200

201 - 250

Tees Mouth

Tees Barrage

TITLE

FIGURE 14B-C2
NFPD FRESHWATER FISH COUNTS
BROWN/SEA TROUT

REFERENCE


NZT_200401_FEB_S_14B-
C2_v2

SHEET NUMBER

1 of 1

DATE

10/06/2020



This drawing has been produced for the use of AECOMs client. It may not be used, modified or relied upon by third parties, except as agreed by AECOM or as required by law. AECOM accepts no responsibility, and denies any liability whatsoever, to any party that uses or relies upon this drawing without AECOMs express written consent. All dimensions are indicative and in metres unless otherwise noted. Do not scale this document.



PROJECT

NET ZERO TEESSIDE

CLIENT

NTZ POWER AND NZNS STORAGE

KEY

NFPD Fish Count Data - European Eel

Total Count

1 - 50

51 - 100

101 - 150

151 - 200

201 - 250

Observed Abundance

1 - 9

10 - 99

100 - 999

1000 - 9999

Tees Mouth

Tees Barrage

TITLE

FIGURE 14B-C3
NFPD FRESHWATER FISH COUNTS
EUROPEAN EEL

REFERENCE


NZT_200401_FEB_S_14B-C3

SHEET NUMBER

1 of 1

DATE

10/06/2020



This drawing has been produced for the use of AECOM client. It may not be used, modified or relied upon by third parties, except as agreed by AECOM or as required by law. AECOM accepts no responsibility, and denies any liability whatsoever, to any party that uses or relies upon this drawing without AECOM's express written consent. All dimensions are indicative and in metres unless otherwise noted. Do not scale this document.



PROJECT

NET ZERO TEESSIDE

CLIENT

NTZ POWER AND NZNS STORAGE

KEY

NFPD Fish Count Data - Lamprey

Total Count

1 - 4

5 - 8

9 - 12

13 - 16

17 - 20

Observed Abundance

1 - 9

10 - 99

100 - 999

1000 - 9999

Tees Mouth

Tees Barrage

TITLE

FIGURE 14B-C4
NFPD FRESHWATER FISH COUNTS
LAMPREY

REFERENCE

NZT_200401_FEB_S_14B-
C4_v2


SHEET NUMBER

1 of 1

DATE

10/06/2020


Figures



This drawing has been produced for the use of AECOM's client. It may not be used, modified or relied upon by third parties, except as agreed by AECOM or as required by law. AECOM accepts no responsibility, and denies any liability whatsoever, to any party that uses or relies upon this drawing without AECOM's express written consent. All dimensions are indicative and in metres unless otherwise noted. Do not scale this document.



PROJECT	
NET ZERO TEESSIDE	
CLIENT	
NZT POWER AND NZNS STORAGE	
KEY	
	10km Study Area
	Site Boundary
	North East Inshore Marine Plan Area (MMO)
	Statistical Rectangle (ICES)
	North Eastern Inshore Fisheries and Conservation Authority District Boundary
TITLE	
FIGURE 14B-1	
STUDY AREA FOR THE FISHERIES AND FISH ECOLOGY BASELINE	
REFERENCE	
NZT_200610_FEB_S_14B-1_v3	
SHEET NUMBER	DATE
1 of 1	10/06/2020



This drawing has been produced for the use of AECOM's client. It may not be used, modified or relied upon by third parties, except as agreed by AECOM or as required by law. AECOM accepts no responsibility, and denies any liability whatsoever, to any party that uses or relies upon this drawing without AECOM's express written consent. All dimensions are indicative and in metres unless otherwise noted. Do not scale this document.



PROJECT
NET ZERO TEESSIDE

CLIENT
NZT POWER AND NZNS STORAGE


- KEY
- 10km Study Area
 - Site Boundary
 - Site of Special Scientific Interest
 - Special Area of Conservation
 - Special Protection Area

TITLE
FIGURE 14B-2
MARINE DESIGNATED SITES WHICH FALL
WITHIN THE STUDY AREA FOR THE
FISHERIES AND FISH ECOLOGY BASELINE

REFERENCE
NZT_200610_FEB_S_14B-2_v3

SHEET NUMBER
1 of 1

DATE
10/06/20



This drawing has been produced for the use of AECOM client. It may not be used, modified or relied upon by third parties, except as agreed by AECOM or as required by law. AECOM accepts no responsibility, and denies any liability whatsoever, to any party that uses or relies upon this drawing without AECOM's express written consent. All dimensions are indicative and in metres unless otherwise noted. Do not scale this document.



PROJECT

NET ZERO TEESSIDE

CLIENT

NZT POWER AND NZNS STORAGE

KEY

- 10km Study Area
- Site Boundary
- Herring Nursery Ground (2010)
- High Intensity
- Low Intensity

TITLE

FIGURE 14B-3a
NURSERY GROUNDS WHICH FALL WITHIN
THE STUDY AREA FOR THE PROPOSED
DEVELOPMENT: HERRING
(ELLIS ET AL., 2012)

REFERENCE


NZT_200610_FEB_S_14B-3a_v3

SHEET NUMBER

1 of 1

DATE


10/06/20



This drawing has been produced for the use of AECOM client. It may not be used, modified or relied upon by third parties, except as agreed by AECOM or as required by law. AECOM accepts no responsibility, and denies any liability whatsoever, to any party that uses or relies upon this drawing without AECOM's express written consent. All dimensions are indicative and in metres unless otherwise noted. Do not scale this document.




PROJECT	NET ZERO TEESSIDE
CLIENT	NZT POWER AND NZNS STORAGE
KEY	<div><div><div></div><div>10km Study Area</div></div><div><div></div><div>Site Boundary</div></div><div><div></div><div>Anglerfish Nursery Ground (2010)</div></div><div><div></div><div>Low Intensity</div></div></div>
TITLE	FIGURE 14B-3b NURSERY GROUNDS WHICH FALL WITHIN THE STUDY AREA FOR THE PROPOSED DEVELOPMENT: ANGLERFISH (ELLIS ET AL., 2012)
REFERENCE	NZT_200610_FEB_S_14B-3b_v3
SHEET NUMBER	1 of 1
DATE	10/06/20



This drawing has been produced for the use of AECOM's client. It may not be used, modified or relied upon by third parties, except as agreed by AECOM or as required by law. AECOM accepts no responsibility, and denies any liability whatsoever, to any party that uses or relies upon this drawing without AECOM's express written consent. All dimensions are indicative and in metres unless otherwise noted. Do not scale this document.




PROJECT	NET ZERO TEESSIDE	
CLIENT	NZT POWER AND NZNS STORAGE	
KEY	<div><div></div>10km Study Area</div> <div><div></div>Site Boundary</div> <div><div></div>Plaice Spawning Ground (2010)</div> <div><div></div>Plaice Nursery Ground (2010)</div> <div><div></div>Low Intensity</div>	
TITLE	FIGURE 14B-3c NURSERY GROUNDS WHICH FALL WITHIN THE STUDY AREA FOR THE PROPOSED DEVELOPMENT: PLAICE (ELLIS ET AL., 2012)	
REFERENCE	NZT_200610_FEB_S_14B-3c_v3	
SHEET NUMBER	1 of 1	DATE 10/06/20



This drawing has been produced for the use of AECOM client. It may not be used, modified or relied upon by third parties, except as agreed by AECOM or as required by law. AECOM accepts no responsibility, and denies any liability whatsoever, to any party that uses or relies upon this drawing without AECOM's express written consent. All dimensions are indicative and in metres unless otherwise noted. Do not scale this document.




PROJECT	NET ZERO TEESSIDE
CLIENT	NZT POWER AND NZNS STORAGE
KEY	<div><div><div></div><div>10km Study Area</div></div><div><div></div><div>Site Boundary</div></div><div><div></div><div>Cod Nursery Ground (2010)</div></div><div><div></div><div>High Intensity</div></div></div>
TITLE	FIGURE 14B-3d NURSERY GROUNDS WHICH FALL WITHIN THE STUDY AREA FOR THE PROPOSED DEVELOPMENT: COD (ELLIS ET AL., 2012)
REFERENCE	NZT_200610_FEB_S_14B-3d_v3
SHEET NUMBER	1 of 1
DATE	10/06/20



This drawing has been produced for the use of AECOM client. It may not be used, modified or relied upon by third parties, except as agreed by AECOM or as required by law. AECOM accepts no responsibility, and denies any liability whatsoever, to any party that uses or relies upon this drawing without AECOM's express written consent. All dimensions are indicative and in metres unless otherwise noted. Do not scale this document.




PROJECT	NET ZERO TEESSIDE
CLIENT	NZT POWER AND NZNS STORAGE
KEY	<div><div><div></div><div>10km Study Area</div></div><div><div></div><div>Site Boundary</div></div><div><div></div><div>Whiting Nursery Ground (2010)</div></div><div><div></div><div>High Intensity</div></div></div>
TITLE	FIGURE 14B-3e NURSERY GROUNDS WHICH FALL WITHIN THE STUDY AREA FOR THE PROPOSED DEVELOPMENT: WHITING (ELLIS ET AL., 2012)
REFERENCE	NZT_200610_FEB_S_14B-3e_v3
SHEET NUMBER	1 of 1
DATE	10/06/20



This drawing has been produced for the use of AECOM client. It may not be used, modified or relied upon by third parties, except as agreed by AECOM or as required by law. AECOM accepts no responsibility, and denies any liability whatsoever, to any party that uses or relies upon this drawing without AECOM's express written consent. All dimensions are indicative and in metres unless otherwise noted. Do not scale this document.



PROJECT	
NET ZERO TEESSIDE	
CLIENT	
NZT POWER AND NZNS STORAGE	
KEY	
	10km Study Area
	Site Boundary
Spurdog Nursery Ground (2010)	
	Low Intensity
TITLE	
FIGURE 14B-3f	
NURSERY GROUNDS WHICH FALL WITHIN	
THE STUDY AREA FOR THE PROPOSED	
DEVELOPMENT: SPURDOG	
(ELLIS ET AL., 2012)	
REFERENCE	
NZT_200610_FEB_S_14B-3f_v3	
SHEET NUMBER	
1 of 1	
DATE	
10/06/20	



PROJECT
NET ZERO TEESIDE

CLIENT
NZT POWER AND NZNS STORAGE

- KEY
- Site Boundary
 - Tees Barrage
 - Tees Mouth
 - TRaC Fish Counts

TITLE
FIGURE 14B-4
TRAC FISH COUNT SAMPLING LOCATIONS
(ENVIRONMENT AGENCY 2019A)

REFERENCE
NZT_200619_FEB_S_14B-4_v3

SHEET NUMBER
1 of 1

DATE
19/06/2020

This drawing has been produced for the use of AECOM's client. It may not be used, modified or relied upon by third parties, except as agreed by AECOM or as required by law. AECOM accepts no responsibility, and denies any liability whatsoever, for any party that uses or relies upon this drawing without AECOM's express written consent. All dimensions are indicative and in metres unless otherwise noted. Do not scale this document.